COHORT PROFILE

Cohort Profile: Wisconsin longitudinal study (WLS)

Pamela Herd,¹* Deborah Carr² and Carol Roan¹

¹University of Wisconsin, Madison, WI, USA and ²Rutgers University, New Brunswick, NJ, USA

*Corresponding author. University of Wisconsin, Madison, 1180 Observatory Drive, Madison, WI 53706, USA. E-mail: pherd@lafollette.wisc.edu

Accepted 9 October 2012

The Wisconsin Longitudinal Study (WLS) is a longitudinal study of men and women who graduated from Wisconsin high schools in 1957 and one of their randomly selected siblings. Wisconsin is located in the upper midwest of the United States and had a population of approximately 14 000 000 in 1957, making it the 14th most populous state at that time. Data spanning almost 60 years allow researchers to link family background, adolescent characteristics, educational experiences, employment experiences, income, wealth, family formation and social and religious engagement to midlife and late-life physical health, mental health, psychological well-being, cognition, end of life planning and mortality. The WLS is one of the few longitudinal data sets that include an administrative measure of cognition from childhood. Further, recently collected saliva samples allow researchers to explore the inter-relationships among genes, behaviours and environment, including genetic determinants of behaviours (e.g. educational attainment); the interactions between genes and environment; and how these interactions predict behaviours. Most panel members were born in 1939, and the sample is broadly representative of White, non-Hispanic American men and women who have completed at least a high school education. Siblings cover several adjoining cohorts: they were born primarily between 1930 and 1948. At each interview, about two-thirds of the sample lived in Wisconsin, and about one-third lived elsewhere in the United States or abroad. The data, along with documentation, are publicly accessible and can be accessed at http://www.ssc.wisc.edu/wlsresearch/. Requests for protected data or assistance should be sent to wls@ssc.wisc.edu.

Why was the cohort set up?

The origins of the Wisconsin Longitudinal Study (WLS) can be traced back to a state-sponsored questionnaire administered during the spring of 1957 to all students in their final year of high school, in all Wisconsin high schools. In the United States, students enter high school around age 14 and typically graduate within 4 years, around age 18. Approximately 75% of students in Wisconsin graduated from high school in the late 1950s.¹ The survey was administered because the state officials wanted to know the demand for post-high school education, whether promising high school students were choosing to attend college or university and what circumstances contributed to their education plans.² In 1964, 7 years after the original question-naire was administered, William H. Sewell, a researcher and Professor of Sociology at the University of Wisconsin-Madison, randomly selected one-third of

the members of the class of 1957 for further data collection. The original goal of the follow-up study was to examine how family background and childhood experiences and aspirations affected educational and occupational attainment. The survey has changed as the participants have aged to cover the relevant aspects of their life course, from family formation and occupational experiences to retirement and health and well-being at older ages.

Who is in the cohort?

WLS is based on a 1/3 random sample of all Wisconsin high school graduates in 1957 (n = 10317) born between 1938 and 1940. Nearly every student in their final year of high school completed the original survey, of whom a random 1/3 sample was drawn for the subsequent surveys. The original questionnaire covered topics such as the student's educational, occupational and marital aspirations, high school coursework and parental and teacher encouragement for post-high school schooling. The sample also includes a randomly selected sibling of these graduates, the majority of whom were born between 1930 and 1948 (n = 7928).^{3,4} Spouses of the graduates and siblings were also interviewed in 2004. Sewell and Hauser⁵ estimated that \sim 75% of Wisconsin youth graduated from high schools in the late 1950s. About 7% of siblings in the WLS did not complete high school, along with 9% of graduate spouses and 12% of siblings' spouses. Despite the educational selectivity of the graduate sample, there is substantial heterogeneity in social origins. For example, Hauser and Sweeney⁶ estimated that >20% of the graduates lived in a household where the 4-year average of parents' income, 1957-1960, fell below the official poverty line. There is only a handful of African American, Hispanic or Asian persons in the WLS, which reflects the very small number of minorities among Wisconsin high school graduates when the WLS began.⁶ Despite these limitations, the WLS provides valuable information about White non-Hispanic cohorts born in 1930s and 1940s. Among Americans aged 60-64 in 2000, 66.7% are non-Hispanic Whites who completed at least 12 years of schooling.⁷

How often have they been followed up?

The graduate respondents originally completed an in-person questionnaire in 1957, which was followed with data collection in 1964 (a mail survey of parents), 1975 (telephone survey), 1993 (telephone and mail surveys), 2004 (telephone and mail surveys) and 2011 (an in-person survey and mail survey—data collection under way). The paired sibling was randomly selected from a roster of all siblings except when the graduate was a twin, in which case the twin was selected. Roughly 2000 siblings were empanelled in 1977, and the full sibling sample was implemented in 1993. Once empanelled, siblings receive the same survey instrument as the graduate. In 2004, spouses of the siblings and graduates completed phone interviews: http://www.ssc.wisc.edu/wlsresearch/documen tation/retention/cor1004 retention.pdf

As Table 1 demonstrates, response rates for the survey, especially for the graduates, have generally been high, especially considering the duration of the panel. Attrition has been due primarily to mortality, and response rates do exclude those who are deceased at each survey round. Generally, as Table 1 shows, response rates for siblings have been lower than response rates for the graduates. Further, response rates in 2004 interviews with spouses, especially for the spouses of the siblings, were low.

Table 2 shows predictors of attrition in WLS, which are similar to other studies, including being male, having lower levels of educational attainment and poorer health.¹ Although the distribution of the sample has changed across time on measures of sex, parental SES, education, IQ and health, Table 2 reveals that the magnitude of these differences is not large. Detailed analyses of non-response for the graduates and siblings show that higher response rates among women and those with higher educational attainment are explained by adolescent cognition, academic performance in high school and participation in civic organizations and volunteering.⁸ Hauser (2005) provides a detailed analysis of nonresponse in the WLS over the history of the survey.⁹

What has been measured?

Table 3 describes the content of the surveys. The survey originally covered key aspects of early life course experiences including family, educational and occupational attainment. By the time the graduates had reached their early 50s, the study shifted towards topics most relevant for midlife and later life, especially health. Table 3 provides only a general overview of content. The 2004 and 2011 surveys, for example, each include around 20000 variables. As with most other surveys, financial questions lead to the most item non-response. To address this non-response, WLS implemented a bracketing technique for questioning reluctant respondents and provides imputed data for earnings, incomes and assets since 1975 for graduates and since 1993 for siblings.

The WLS also now includes genetic data obtained from saliva samples collected from respondents in 2006–2007 using Oragene kits and a mailback protocol patterned closely on a previous Swedish study.¹⁰ Oragene kits were selected because of their ability to be used in a mailback protocol (e.g. no need for immediate freezing) and their high average DNA yield relative to widely used alternatives for mailback

		Gradua	ate sample		Sibling sample				
	Sample frame	Deceased	Completed interviews	Response rate	Sample frame ^a	Deceased	Completed interviews	Response rate	
1957	10317		10317	100%					
1964									
Mail	10317	71	8922	90%					
1975									
Mail	10317	175	9138	92%	2438	9	2133	87.81%	
1993									
Phone	10317	587	8493	86%	7812	462	4804	65.36%	
Mail	10317	587	6845	69%	7812	462	4065	55.31%	
Either phone or mail	10317	587	8493	86%	7812	462	5365	72.99%	
2004									
Phone	10317	1288	7265	73%	7928	1226	4298	64.13%	
Mail	10317	1288	6845	69%	7928	1226	4004	59.74%	
Either phone or mail	10317	1288	7732	78%	7928	1226	4703	70.17%	
2004 spouse	7483	22	4076	58%	5750	22	2180	38.06%	

Table 1 Response rates in the WLS

^aThe sample frame for siblings for 1975 was a 1/3 random sample of one random living sibling for each graduate with a sibling. The 1993 eligible sibling sample excluded: siblings who were in the 1975 sample frame, but were not interviewed; siblings of graduates not interviewed in 1975; and interviews not completed owing to lacking funds. The 2004 eligible siblings sample excluded: siblings sample frame, but were not interviewed; siblings of graduates not interviewed in 1975 and/or 1993 sample frame, but were not interviewed; siblings of graduates not interviewed in 1975 or 1993.

protocols available at the time. Additional samples are currently being collected in-person from those who did not provide a sample in 2006–2007.

Finally, since its inception the WLS has collected a range of non-survey-derived measures, which are detailed in Table 4. As noted in Table 4, data users may access participants' Medicare claims records. Medicare is the universal federally funded health insurance program that covers nearly everyone in the United States age 65 and older. Data users will also have access to participants' Social Security lifetime earnings and benefits records. Social Security is the federal government's old-age public pension program that covers nearly all Americans. Access to Medicare and Social Security data, however, will require measures added to the downloadable public data given the sensitivity of these data. These data will be available in 2013. Information about access can be obtained via emails to wls@ssc.wisc.edu.

What are the main strengths and weaknesses?

Strengths

The WLS is one of the longest running studies of a single cohort of adults in the United States, spanning from participants' senior year in high school in 1957 to 2011 with our ongoing data collection. Sample members are tracked through to their deaths.

Further, the graduate data are enhanced by parallel survey data on one randomly selected sibling per graduate, enabling researchers to explore sibling similarities in life course experiences. In 2004, spouses of the graduates and siblings also completed interviews, thus the WLS is well-suited for couple-level analyses. Retention is remarkably high given the length of the panel.

The WLS is a unique data resource, when compared with other large population-based longitudinal studies in the United States, owing to its duration, sibling data and extensive family histories and background. Survey content encompasses nearly every facet of the participants' lives. The WLS has been touted as the most comprehensive long-standing cohort study cur-rently in existence.¹¹ Some elements of the survey that are particularly distinctive relative to other existing longitudinal cohort studies include administrative data on the participants' (including spouses) IQ scores when in high school, and full lifetime employment histories (including job characteristics). Most recently, WLS obtained high school yearbooks for the graduates. Researchers coded the yearbook data for a range of indicators including youthful facial attractiveness, body weight and school activity profiles, allowing one-of-a-kind analyses of the long-term influences of early life characteristics.

The WLS' distinctive attributes will be especially valuable when examined in tandem with the recently added genetic data. The length of the panel presents

	Graduate				Sibling		Spouse				
	1957	1975	1993	2004	1977	1993	2004	Grad report of education 2		Interviewed spouses	
Graduate's educational a	ttainmen	t in 197	5					Grad report of	spouse's	education	
<high school<="" td=""><td></td><td></td><td></td><td></td><td>7.64</td><td>6.53</td><td>4.81</td><td>7.81</td><td></td><td>5.42</td></high>					7.64	6.53	4.81	7.81		5.42	
HS only		60.05	59.08	57.29	52.87	48.05	44.8	53.56		50.58	
>High school		39.95	40.93	42.71	39.49	45.41	50.39	38.63		44	
Self-rated health								Grad report of spouse's health			
Very poor			0.44	0.29		0.45	0.29	35.66		37.58	
Poor			1.25	0.88		1.78	1.15	47.43		47.1	
Fair			9.9	9.08		11.09	9.68	12.5		11.74	
Good			59.59	59.93		59.92	60.18	3.56		2.9	
Excellent			28.83	29.81		26.76	28.7	0.85		0.67	
Sex											
Male	48.38	47.38	46.86	46.29	49.15	48.47	47.59				
Female	51.62	52.62	53.14	51.62	50.85	51.53	52.41				
Father's education											
<high school<="" td=""><td>61.36</td><td>61.72</td><td>61.55</td><td>60.66</td><td>61.84</td><td>60.49</td><td>59.25</td><td></td><td></td><td></td></high>	61.36	61.72	61.55	60.66	61.84	60.49	59.25				
HS degree	23.81	24.18	24.2	24.38	24.5	24.52	24.69				
>High school	14.83	14.11	14.25	14.96	13.67	14.99	16.06				
Mother's education											
<high school<="" td=""><td>49.31</td><td>49.41</td><td>48.85</td><td>48.13</td><td>48.83</td><td>48.34</td><td>46.09</td><td></td><td></td><td></td></high>	49.31	49.41	48.85	48.13	48.83	48.34	46.09				
HS degree	36.41	36.65	36.86	37.05	36.94	37.36	37.87				
>High school	14.28	13.94	14.29	14.81	14.23	14.31	16.04				
Mean Henmon–Nelson standardized IQ score	100.459	100.79	101.19	101.92	101.93	102.89	104.43				

Table 2 Major correlates of attrition in the WLS

Grad, graduate respondent.

researchers the opportunity to look at links between genes and phenotypic data for many participants from early adolescence into their early 70s. The WLS offers extensive measures of the environment in a 55-year longitudinal framework, allowing for attention to life course processes. This nearly full life course longitudinal panel, with extensive family, employment, educational, psychological, cognitive and health data, will provide a unique opportunity to evaluate interactions between gene variants and a series of carefully measured prospective variables. Further, WLS is unique as a population-based study of this size and breadth that includes genetic data on siblings.

Weaknesses

Despite these many strengths, the WLS has several weaknesses pertaining primarily to its limited generalizability. By design, all members of the graduate sample are high school graduates, and nearly all are White—reflecting the composition of Wisconsin high school graduates in the late 1950s. Thus, the findings necessarily apply to a single cohort of White adults with at least a high school diploma (although the sibling and spouse samples show greater educational and age heterogeneity). Despite these limitations, the sample is broadly representative of older White American men and women who have completed at least a high school education. Non-Hispanic Whites who have completed at least a high school education accounted for more than two-thirds all American women and men ages 70–74 in the 2010 US Census.

However, sample homogeneity provides certain advantages. First, sample homogeneity has strengthened many studies by helping to rule out unobserved variable bias as a source of causal relationships. Notable studies include Snowdon's sample of nuns and the Framingham Study.^{12,13} Indeed, most multivariate statistical analyses are intended to reduce sample heterogeneity. Second, sample homogeneity is particularly useful for analyses using the genetic data. These features can strengthen the power of genetic studies using the WLS against population stratification and other factors.¹⁴

A further limitation is that detailed physical and mental health data were first obtained in the 1993

38 INTERNATIONAL JOURNAL OF EPIDEMIOLOGY

Table 3 Summary of WLS content	Table 3 Continued					
1957 (Self-administered questionnaire)	End of life preparations					
Marital aspirations and plans	End of life experiences (death of spouse or parent)					
Occupational aspirations and plans	Menstruation, menopause and hysterectomy					
High school coursework	Hostility, anger, anxiety					
Friends' occupational aspirations and plans	Masculinity scale					
Parents' occupational and educational attainment	Social support and relationships					
1964 (Parent reporting on the graduate)	Marital quality					
Detailed current schooling and attainment since high school	Health Utility Index (functional measure) More detailed measures of chronic illnesses than collected					
Current and past job history	in 1992					
Marital status	Cognition-letter/category fluency					
Spouse's current schooling and occupation	Cognition-similarities					
1975 (Graduate and sibling respondents)	Cogntion-word recall					
1964 Measures	Health care access and utilization					
Civic engagement (i.e. participation in fraternal	Retrospective childhood health ^a					
organizations)	Retrospective childhood abuse ^a					
Time spent with family and friends	2011 (Graduate and sibling respondents)					
Religious orientation and participation	2004 Measures repeated					
Marital history	Network data linking study participants					
Detailed military history	Financial literacy					
Detailed unemployment history	Retirement attitudes					
Current/last job characteristics and aspirations	Risk preferences					
Graduate and spouses' earnings	Questions regarding the great recession					
Fertility history (number, sex, birth year)	Special survey of parents of children with developmental					
Educational, occupational and income aspirations and expectations for a selected child	disabilities or significant mental illness Activities of daily living/instrumental activities of daily					
1992 (Graduate and sibling respondents)	living					
1975 Measures repeated	Cognition-delayed and immediate recall					
Detailed assets summary (including intertransfers)	Cogntion-cookie theft description					
Detailed income summary	Extensive cognitive testing for 10% of the sample					
Pension summary	Waist-hip ratio					
Future plans and retirement	Peak-flow capacity					
Health insurance information	Time gait test					
Caregiving	Chair rise					
Selected child (education, employment, marital status, relationship with parents)	Vision screener Medication inventory					
Chronic conditions	Medication inventory Detailed questions on diabetes knowledge					
Physical symptoms						
Physical limitations/functional limitations	Anaesthesia experiences					
Menopause and hormone therapy	Elder abuse					
Smoking	wave, when the respondents were ages 53–54, thus					
2004 (Graduate and sibling respondents and their spouses)	limiting researchers' ability to track health over the					
1992 Measures repeated	full life course. This limitation was partially corrected by asking participants in 2004 to retrospectively					
Network data linking study participants	evaluate their physical and mental health at age 16					
Volunteering	Further, the WLS collects complete employment his- tories, including spells of non-employment due to dis-					
Internet use	ability Thus researchers can construct broad provie					

ability. Thus, researchers can construct broad proxies

for poor health during young adulthood.

Table 3 Continued

Downloaded from https://academic.oup.com/ije/article-abstract/43/1/34/729687/Cohort-Profile-Wisconsin-longitudinal-study-WLS by University of Wisconsin-Madison user on 21 September 2017

(continued)

Table 4 WLS adminstrative data

Mortality data (updated quarterly)

Cause of death data (updated annually)

Wisconsin state tumour registry data

Medicare records

Social security lifetime earnings/benefits records

Geographical identifiers

Graduates' parents' earnings 1957-1960

Henmon-Nelson IQ scores 1954-1957

High school rank

School district data (teacher/student ratios, funding)

College/university characteristics

Data from high school yearbooks

Facial attractiveness

Facial mass index (obseity)

Participation in school activities (sports and interest clubs)

Respondent's voting data

Area Resource Files

6000 variables containing information on health care facilities, health professions, measures of resource scarcity, health status, economic activity, health training programs and socio-economic and environmental characteristics

What has it found? Key findings and publications

The WLS has been regarded historically as the premier data set for exploring status attainment processes over the life course. In recent decades, WLS data collection has focused on health and aging, thus generating important publications exploring three main areas: the long-term influences of early life factors on adult health; the use of sibling models to address the role of early life experiences; and the social stratification of healthy aging. Further, recent work incorporating genetic data points to the enormous potential of WLS to contribute to genetic research. A complete and searchable bibliography of publications is available at http://www.ssc.wisc.edu/wlsresearch/publications/.

First, the collection of data over a >50-year span has positioned the WLS as a premier data source for exploring the long-term physical and mental health consequences of a range of adolescent and young adulthood characteristics and experiences. The study has demonstrated how adversities in childhood and early life shape health well into late life. For example, negative childhood experiences, especially childhood physical abuse, exert strong and negative impacts on health in late life.^{15,16} Being raised in a poor family, lower levels of educational and occupational attainment, in addition to working in jobs that lack autonomy and control, predict a range of poor health and mortality outcomes in mid to late life, including an increased risk of breast cancer mortality.^{17–20} Extensive social measures in the study collected throughout the participants' entire lives have also been used to show that volunteering and positive relationships with parents, children and friends all exert strong and positive influences on health in late life.^{21,22} Finally, the WLS data have more recently contributed to debates in cognitive epidemiology focused on how early life cognition influences late life health and mortality. In particular, these studies found that one's academic performance in high school was a much stronger predictor of late life health and mortality than was cognition.²³ Further, measures of cognition, from early to late life, and detailed life course measures of alcohol use, helped to identify the causal ordering of the association between cognition and drinking. Analysis shows that those with higher IQs in adolescence consume more alcohol than do those with lower IQs.²

Second, the sibling data have helped clarify the role that early life family context plays in a range of late life outcomes. Sibling models enable researchers to control for unobserved heterogeneity that results from shared early childhood experiences as well as focus on sibling resemblance. The WLS sibling data have been used to demonstrate that the effect of family background on occupational attainment is fully accounted for by its impact on cognition and educational attainment, as opposed to any kind of direct effect of family of origin.²⁵ More recent work focused on health outcomes demonstrated that although siblings exert important influences on each other's childhood development, they do not exert any influence on each other's health in later life.²⁶ Further, WLS sibling models revealed that childhood family context could not explain links between IQ and health in late life. Prior work in this area was limited by its inability to evaluate whether the link between IQ and health was a product of early life family experiences. The current collection of DNA data will allow investigators to further explore the social and genetic bases of sibling resemblance in future studies.

Third, the breadth and depth of measures obtained in the most recent waves of the WLS have generated a range of publications on social inequities in a range of later-life health experiences, health experiences not often measured in comparable cohort studies. For example, the WLS' extensive range of measures on women's reproductive health led researchers to discover that higher incomes and educational attainment reduce the risk of hysterectomy owing to health insurance advantages associated with better jobs.²⁷ Higher levels of educational attainment, occupational attainment and cognition were positively correlated with age of menstruation cessation.²⁸ Although higher socio-economic status is almost uniformly linked to positive health outcomes, analyses of WLS data revealed that higher SES is associated with elevated risk of breast cancer, which may be due to women with higher SES tending to use hormone replacement therapy.²⁹ Further, the (on average) older age of women with higher SES at first birth also partly contributes to their elevated risk of breast cancer.²⁰

The WLS has also shown how SES affects medical care. Those with higher levels of educational attainment have made more successful use of the internet to obtain health-related information,³⁰ and participate more extensively in health care decision making.³¹ Higher educational attainment affects how people plan for the end of their lives. Those with higher educational attainment and richer economic assets are more likely to engage in advance care planning (i.e. living wills and informal discussion with family members).³² Those with higher educational attainment were also more likely to want to continue medical treatments even if faced with the hypothetical scenario of terminal illness with physical pain.³³

Finally, the recent inclusion of genetic data to the WLS has pointed to the role the study can play in the growing body of genetic research. In addition to its wealth of life course phenotypic data, the WLS' advantages include its relatively large sample size, which helps address the common problem of statistically underpowered analyses that have frequently led to false findings. One recent example of how a larger sample can address false findings is a study by Christopher Chabris et al.³⁴ who used administrative measures of intelligence from adolescence (the Henmon-Nelson intelligence test) and genotypes for 13 of the most common single nucleotide polymorphisms (the most frequent form of genetic variation found among people) reported to be associated with intelligence in existing published studies. They found that most of these associations were likely to be false positives and will not consistently replicate in other samples.³

In sum, the WLS data reveal the complex ways in which social statuses, roles and experiences over the life course affect midlife and late life health, with attention to the psychosocial, cognitive and biological pathways that contribute to health inequalities.

Can I get hold of the data? Where can I find out more?

WLS data collection has been funded by the United States' National Institute on Aging since 1991. Detailed documentation and data can be down-loaded for free at http://www.ssc.wisc.edu/wlsre search/. Protected measures, which are those measures that increase a subject's risk of identifiability, are available to qualified researchers by request. Researchers wanting to use the genetic data may submit a proposal that will be reviewed by an independent board. The board reviews the proposals only to ensure that the confidentiality of participants' information will be protected. To receive more information on how to access protected measures (including the genetic data) please email wls@ssc.wisc.edu.

Funding

Since 1991, the WLS has been supported principally by the National Institutes for Health, National Institute on Aging, with additional support from the Vilas Estate Trust, the National Science Foundation, the Spencer Foundation, and the Graduate School of the University of Wisconsin-Madison.

Conflict of interest: None declared.

KEY MESSAGES

- A nearly 60-year longitudinal panel cohort design has demonstrated how adversities in childhood and early life, including childhood physical abuse and limited economic, cognitive and social resources, exert strong and negative impacts on health in late life.
- The sibling panel design has helped demonstrate the role that one's family plays in a range of late life outcomes, while simultaneously accounting for unobserved heterogeneity that results from shared early childhood experiences.
- The breadth and depth of measures obtained in the most recent waves of the WLS have demonstrated how low educational attainment, income and occupational attainment increase the risk for negative health outcomes and medical care experiences, such as general health and mortality, cardiovascular mortality, hysterectomy and the early cessation of menstruation. Studies have also demonstrated how individuals with more education and income engage more effectively in health care decision making and end of life care planning and decisions.
- The large sample size and rich array of life course phenotypic data, combined with the recent addition of genetic data, has begun to allow researchers to explore the interrelationships among genes, behaviours and environment.

References

- ¹ Hauser RM, Willis RJ. Survey design and methodology in the Health and Retirement Study and the Wisconsin Longitudinal Study. In: Waite LJ (ed.). *Aging, Health, and Public Policy: Demographic and Economic Perspectives*. New York: Population Council, 2005, pp. 209–35.
- ² Little James K. A State-Wide Inquiry into Decisions of Youth about Education beyond High School. Madison, WI: University of Wisconsin Madison, 1958.
- ³ Sewell WH, Hauser RM, Springer KW, Hauser TS. As we age: the Wisconsin Longitudinal Study. In: Leicht K (ed.). *Research in Social Stratification and Mobility*. London: Elsevier, 2004, pp. 3–111.
- ⁴ Hauser R. Causes and consequences of cognitive functioning across the life course. *Educ Res* 2009;**39**:95–109.
- ⁵ Sewell WH, Hauser RM. *Education, Occupation, and Earnings: Achievement in the Early Career*. New York: Academic Press, 1975.
- ⁶ Hauser RM, Sweeney MM. Does adolescent poverty affect the life chances of high school graduates? In: Duncan G, Brooks-Gunn J (eds) *Growing up Poor*. New York: Russell Sage Foundation, 1997, pp. 541–95.
- ⁷ U.S. Bureau of the Census. *Educational Attainment in the United States: March 2000*. Washington, DC: Government Printing Office, 2000.
- ⁸ Freese J. Cognitive Skills and Survey Nonresponse: Evidence from Two Longitudinal Studies. Madison, WI: Center for Demography and Ecology, University of Wisconsin, 2006. CDE Working Paper No. 2006–10.
- ⁹ Hauser RM. Survey response in the long run: the Wisconsin longitudinal study. *Field Methods* 2005;**17**:3–29.
- ¹⁰ Rylander-Rudqvist T, Håkansson N, Tybring G, Wolk A. Quality and quantity of saliva DNA obtained from the self-administrated Oragene method. *Cancer Epidemiol Biomarkers Prev* 2006;**15**:1742–45.
- ¹¹ Singer BH, Ryff CD (eds). National Research Council. New Horizons in Health: An Integrative Approach. Washington, DC: National Academy Press, 2001, p. 105.
- ¹² Snowdon DA. Healthy aging and dementia: findings from the Nun Study. *Ann Int Med* 2003;**139**:450–54.
- ¹³ Levy D, Brink D. A Change of Heart: How the Framingham Heart Study Helped Unravel the Mysteries of Cardiovascular Disease. New York: Knopf, 2005.
- ¹⁴ Ott J, Kamatani Y, Lathrop M. Family-based designs for genome-wide association studies. *Nat Rev Genetics* 2011; 12:465–74.
- ¹⁵ Springer KW, Sheridan J, Kuo D, Carnes M. The long-term health outcomes of childhood abuse: an overview and a call to action. J Gen Int Med 2003;18:864–70.
- ¹⁶ Springer KW, Sheridan J, Kuo D, Carnes M. Long-term physical and mental health consequences of childhood physical abuse: results from a large population-based sample of men and women. *Child Abuse Negl* 2007;**31**: 517–30.
- ¹⁷ Carr D. Socioeconomic background and midlife health in the United States. In: Crystal S, Shea D (eds). *Annual Review of Gerontology and Geriatrics, Volume 22: Economic*

Outcomes in Later Life: Public Policy, Health, and Cumulative Advantage. New York: Springer, 2002, pp. 155–83.

- ¹⁸ Warren JR, Hoonakker P, Carayon P, Brand J. Job characteristics as mediators in SES-health relationships. *Soc Sci Med* 2004;**59:**1367–78.
- ¹⁹ Warren JR, Kuo H-H. How to measure "what people do for a living" in research on the socioeconomic correlates of health. *Ann Epidemiol* 2003;**13**:325–34.
- ²⁰ Pudrovska T, Benedicta A. The role of early-life socioeconomic status in breast cancer incidence and mortality: unraveling life course mechanisms. *J Aging Health* 2012; 24:323–44.
- ²¹ Konrath FA, Lou A, Brown S. Motives for volunteering are associated with mortality risk in older adults. *Health Psychol* 2012;**31**:87–96.
- ²² Seeman TE, Singer BH, Ryff CD, Love GD, Levy-Storms L. Social relationships, gender, and allostatic load across two age cohorts. *Psychosom Med* 2002;64:395–406.
- ²³ Herd P. Education and health in late-life among high school graduates: cognitive versus psychological aspects of human capital. *J Health Soc Behav* 2010;**51**:478–96.
- ²⁴ Krahn D, Freese J, Hauser R, Barry K, Goldman B. Alcohol use and cognition at mid-life: the importance of adjusting for baseline cognitive ability and educational attainment. *Alcohol Clin Exp Res* 2003;**27**:1162–66.
- ²⁵ Hauser RM, Sheridan J, Warren JR. Socioeconomic achievements of siblings in the life course: new findings from the Wisconsin Longitudinal Study. *Res Aging* 1999; 21:338–78.
- ²⁶ Falbo T, Kim S, Chen K. Alternate models of sibling status effects on health in later life. *Dev Psychol* 2009;45: 677–87.
- ²⁷ Marks NF, Shinberg DS. Socioeconomic differences in hysterectomy: the Wisconsin Longitudinal Study. Am J Public Health 1997;87:1507–14.
- ²⁸ Shinberg DS. An event history analysis of age at last menstrual period: correlates of natural and surgical menopause among midlife Wisconsin women. *Soc Sci Med* 1998;**46**:1381–96.
- ²⁹ Marks NF, Shinberg DS. Socioeconomic status differences in hormone therapy. *Am J Epidemiol* 1998;**148**:581–93.
- ³⁰ Flynn KE, Smith MA, Freese J. When do older adults turn to the internet for health information?: findings from the Wisconsin Longitudinal Study. *J Gen Intern Med* 2006;**21**:1295–301.
- ³¹ Flynn KE, Smith MA, Vanness D. A typology of preferences for participation in healthcare decision making. *Soc Sci Med* 2006;**63**:1158–69.
- ³² Carr D, Khodyakov D. End of life health care planning among young-old adults: an assessment of psychosocial influences. J Gerontol B Psychol Sci Soc Sci 2007;62:135–41.
- ³³ Carr D, Moorman S. End-of-life treatment preferences among older adults: an assessment of psychosocial influences. *Sociol Forum (Randolph NJ)* 2009;**24**:754–78.
- ³⁴ Chabris CF, Hebert BM, Benjamin DJ *et al.* Most published genetic associations with general intelligence are probably false positives. *Psychol Sci* 2012;**23**:1314–23.