

Econometrics 710  
Midterm Exam  
March 23, 2000

1. The model is

$$y_i = x_i\beta + e_i \quad E(e_i | x_i) = 0$$

where  $x_i$ ,  $\beta$  and  $e_i$  are scalar. We consider the estimator

$$\tilde{\beta} = \frac{\bar{y}}{\bar{x}} = \frac{\sum_{i=1}^n y_i}{\sum_{i=1}^n x_i}.$$

We assume that  $x_i$  and  $e_i$  have finite fourth moments and that  $\{y_i, x_i\}$  are a random sample (iid).

- Find  $E(\tilde{\beta} | X)$ .
  - Find  $Var(\tilde{\beta} | X)$ .
  - Show that  $\tilde{\beta} \rightarrow_p \beta$  as  $n \rightarrow \infty$ . Does this require any additional assumptions?
  - Find the asymptotic distribution of  $\sqrt{n}(\tilde{\beta} - \beta)$  as  $n \rightarrow \infty$ .
  - Without imposing any additional assumptions, is  $\tilde{\beta}$  necessarily less efficient than OLS? (By efficiency, I mean lower asymptotic variance.)
2. Take the linear regression  $Y = X\beta + e$  with  $E(e_i | x_i) = 0$ . Let  $\theta = 1/\beta_1$  where  $\beta_1$  is the first element of  $\beta$ . Let  $\hat{\beta}$  be the OLS estimator of  $\beta$  and  $\hat{V}$  be the estimator of  $Var(\hat{\beta})$ . Find an asymptotically valid 95% confidence interval for  $\theta$ . (Give the explicit formula as a function of  $\hat{\beta}$  and  $\hat{V}$ .)
3. In the linear regression  $Y = X\beta + e$  with  $E(e_i | x_i) = 0$ , it is known that the true  $\beta$  satisfies the restriction

$$R\beta = 0$$

where  $R$  is a  $q \times k$  matrix with  $q < k$ . Consider the estimator

$$\tilde{\beta} = \hat{\beta} - (X'X)^{-1} R' [R(X'X)^{-1} R']^{-1} R\hat{\beta}.$$

- Show that  $R\tilde{\beta} = 0$ .
- Find  $E(\tilde{\beta} | X)$ .
- Find  $Var(\tilde{\beta} | X)$ . [Hint: First write  $\tilde{\beta}$  as a linear function of  $\hat{\beta}$ .]

- (d) Give an expression for a valid standard error for the elements of  $\tilde{\beta}$ . You do not need to give a proof of validity.
4. Take the linear regression  $Y = X\beta + e$  with  $E(e_i | x_i) = 0$ . For one particular value of  $x$ , the object of interest is the conditional mean

$$E(y_i | x_i = x) = g(x).$$

Describe how you would use the percentile-t bootstrap to construct a confidence interval for  $g(x)$ .