

Econometrics 710
Midterm Exam
March 6, 2003

1. The model is

$$\begin{aligned}y_i &= x_i' \beta + e_i \\ E(x_i e_i) &= 0.\end{aligned}$$

The *ridge regression* estimator for β is

$$\hat{\beta} = (X'X + \lambda I_k)^{-1} (X'Y).$$

Suppose that $\lambda = cn$ with c fixed as $n \rightarrow \infty$. Find the probability limit of $\hat{\beta}$ as $n \rightarrow \infty$.

2. The model is

$$\begin{aligned}y_i &= x_i' \beta + e_i \\ E(x_i e_i) &= 0 \\ \Omega &= E(x_i x_i' e_i^2).\end{aligned}$$

- (a) Find the method of moments estimators $(\hat{\beta}, \hat{\Omega})$ for (β, Ω) .
(b) In this model, are $(\hat{\beta}, \hat{\Omega})$ efficient estimators of (β, Ω) ?
(c) If so, in what sense are they efficient?
3. Suppose we have an estimate $\hat{\beta}$ of $\beta \in R$ such that $\sqrt{n}(\hat{\beta} - \beta) \rightarrow^d N(0, V)$ as $n \rightarrow \infty$, we have a consistent estimator \hat{V} of V , and the parameter of interest is $\theta = \beta^2$.
- (a) Find the asymptotic distribution of $\hat{\theta} = \hat{\beta}^2$.
(b) Use result (a) to form a confidence interval for θ .
(c) What are the consequences if $\beta = 0$?

4. The model is

$$\begin{aligned}y_i &= x_i \beta + e_i \\ E(e_i | x_i) &= 0\end{aligned}$$

where $x_i \in R$. Consider the two estimators

$$\begin{aligned}\hat{\beta} &= \frac{\sum_{i=1}^n x_i y_i}{\sum_{i=1}^n x_i^2} \\ \tilde{\beta} &= \frac{1}{n} \sum_{i=1}^n \frac{y_i}{x_i}.\end{aligned}$$

- (a) Under the stated assumptions, are both estimators consistent for β ?
(b) Are there conditions under which either estimator is efficient?