

ACCUMULATED PREDICTION ERRORS, INFORMATION CRITERIA AND OPTIMAL FORECASTING FOR AUTOREGRESSIVE TIME SERIES

Ching-Kang Ing*
Academia Sinica and National Taiwan University

Abstract

The predictive capability of a modification of Rissanen's accumulated prediction error (APE) criterion, APE_{δ_n} , is investigated in infinite-order autoregressive (AR(∞)) models. Instead of accumulating squares of sequential prediction errors from the beginning, APE_{δ_n} is obtained by summing these squared errors from stage $n\delta_n$, where n is the sample size and $1/n \leq \delta_n \leq 1 - (1/n)$ may depend on n . Under certain regularity conditions, an asymptotic expression is derived for the mean-squared prediction error (MSPE) of an AR predictor with order determined by APE_{δ_n} . This expression shows that the prediction performances of APE_{δ_n} can vary dramatically depending on the choice of δ_n . Another interesting finding is that when δ_n approaches 1 at a certain rate, APE_{δ_n} can achieve asymptotic efficiency in most practical situations. An asymptotic equivalence between APE_{δ_n} and an information criterion with a suitable penalty term is also established from the MSPE point of view. This offers new perspectives for understanding the information- and prediction-based model selection criteria. Finally, we provide the first asymptotic efficiency result for the case when the underlying AR(∞) model is allowed to degenerate to a finite autoregression.

AMS 2000 subject classifications. Primary 60M20; secondary 62F12, 62M10.

Key words and phrases: Accumulated prediction errors, Asymptotic equivalence, Asymptotic efficiency, Information criterion, Order selection, Optimal forecasting

*Institute of Statistical Science, Academia Sinica, 128, Academia Rd. Sec. 2, Taipei 115, Taiwan, R.O.C. E-mail: cking@stat.sinica.edu.tw

1. INTRODUCTION

In the past two decades, investigations on the accumulated prediction error (APE) (Rissanen, 1986) and its variations have attracted considerable attention among researchers from various disciplines. Prior to the early 1990s, a large number of studies focused on its consistency in selecting regression or time series models (e.g., Wax, 1988, Hannan et al., 1989, Hemerly and Davis, 1989, Wei, 1992, Speed and Yu, 1993). However, since proving consistency requires assuming that the true model is included among the family of candidate models (which is rather difficult to justify in practice), recent research has focused more on understanding its statistical properties under possible model misspecification (e.g., Kavalieris, 1989, Wei, 1992, West, 1996, McCracken, 2000, Findley, 2005, Inoue and Kilian, 2006, among others). While a much deeper understanding of the APE in cases of a misspecified model has been gained from these recent efforts, the APE's prediction performance after model selection still remains unclear. This motivated the present study.

To select a model for the realization of a stationary time series, it is common to assume that the realization comes from an autoregressive moving-average (ARMA) process whose AR and MA orders are known to lie within prescribed finite intervals. Then a model selection procedure is used to select orders within these intervals and thereby determine a model for the data. However, as pointed out by Shibata (1980), Goldenshluger and Zeevi (2001) and Ing and Wei (2005), this assumption can rarely be justified in practice, and the less stringent assumption is that the time series data are observations from a linear stationary process. Following this idea, it is assumed in the sequel that observations x_1, \dots, x_n are generated by an $\text{AR}(\infty)$ process $\{x_t\}$, where

$$x_t + \sum_{i=1}^{\infty} a_i x_{t-i} = e_t, t = 0, \pm 1, \pm 2, \dots, \quad (1.1)$$

with the characteristic polynomial $A(z) = 1 + \sum_{i=1}^{\infty} a_i z^i \neq 0$ for all $|z| \leq 1$ and $\{e_t\}$ being a sequence of independent random noises satisfying $E(e_t) = 0$ and $E(e_t^2) = \sigma^2$ for all t . To predict future observations, we consider a family of approximation models $\{\text{AR}(1), \dots, \text{AR}(K_n)\}$, where the maximal order K_n is allowed to tend to ∞ as n does in order to reduce approximation errors. In this framework, the APE value of model $\text{AR}(k)$, $1 \leq k \leq K_n$, is given by

$$\text{APE}(k) = \sum_{i=m}^{n-1} (x_{i+1} - \hat{x}_{i+1}(k))^2, \quad (1.2)$$

where $\hat{x}_{i+1}(k) = -\mathbf{x}'_i(k)\hat{\mathbf{a}}_i(k)$, $\mathbf{x}_i(k) = (x_i, \dots, x_{i-k+1})'$, $\hat{\mathbf{a}}_i(k)$ satisfies

$$-\hat{R}_i(k)\hat{\mathbf{a}}_i(k) = \frac{1}{i - K_n} \sum_{j=K_n}^{i-1} \mathbf{x}_j(k)x_{j+1}, \quad (1.3)$$

with

$$\hat{R}_i(k) = \frac{1}{i - K_n} \sum_{j=K_n}^{i-1} \mathbf{x}_j(k)\mathbf{x}'_j(k), \quad (1.4)$$

and $m \geq K_n + 1$ is the first integer j such that $\hat{\mathbf{a}}_j(K_n)$ is uniquely defined. As observed, $\text{APE}(k)$ measures the performance of $\text{AR}(k)$ when it is used for sequential predictions. Recently, a modification of the APE,

$$\text{APE}_{\delta_n}(k) = \sum_{i=n\delta_n}^{n-1} (x_{i+1} - \hat{x}_{i+1}(k))^2, \quad (1.5)$$

with $1/n \leq \delta_n \leq 1 - (1/n)$ depending on n , has also been considered by several authors, e.g., West (1996), McCracken (2000), and Inoue and Kilian (2006). Since the APE_{δ_n} includes the original APE as a special case, this paper focuses on the APE_{δ_n} . As will be shown later, the performance of the APE_{δ_n} can vary dramatically depending on the choice of δ_n .

In view of (1.5), it is natural to predict the next observation x_{n+1} using $\hat{x}_{n+1}(\hat{k}_{n,\delta_n})$, where

$$\hat{k}_{n,\delta_n} = \arg \min_{1 \leq k \leq K_n} \text{APE}_{\delta_n}(k). \quad (1.6)$$

This type of prediction, targeting future values of the observed time series, is referred to as a *same-realization prediction*. On the other hand, if the process used in estimation (or model selection) and that for prediction are independent, then it is called an *independent-realization prediction* (see Shibata, 1980, Bhansali, 1986, Karagrigoriou, 1997, and Schorfheide, 2005). For differences between these two types of predictions in various time series models, see Kunitomo and Yamamoto (1985), Ing (2001, 2003) and Ing and Wei (2003, 2005). The prediction performance of the APE_{δ_n} after order selection is assessed using the mean-squared prediction error (MSPE) $q_n(\hat{k}_{n,\delta_n})$, where, with $\hat{k}_n = \hat{k}_n(x_1, \dots, x_n) \in \{1, 2, \dots, K_n\}$,

$$q_n(\hat{k}_n) = E \left(x_{n+1} - \hat{x}_{n+1}(\hat{k}_n) \right)^2. \quad (1.7)$$

There are three interrelated issues addressed in this paper. The first one focuses on the asymptotic expression for $q_n(\hat{k}_{n,\delta_n})$. To deal with this problem, in Proposition

2 (see Section 2), we establish a general theory that provides sufficient conditions under which $q_n(\hat{k}_n) - \sigma^2$ can be asymptotically expressed as a sum of two terms that measure the model complexity and the goodness of fit. This result is then applied to the case $\hat{k}_n = \hat{k}_{n,\delta_n}$ with δ_n bounded away from 1; see Theorem 1 in Section 3. A series of examples is given after Theorem 1 to illustrate its implications. In particular, it is shown in Example 1 that when the AR coefficients $\{a_i\}$ decay exponentially (which includes, but is not limited to, the ARMA(p, q) model with $q > 0$ as a special case) and δ_n satisfies $\log \delta_n^{-1} = o(\log n)$, the APE_{δ_n} is asymptotically efficient in the sense of (2.3). However, if the $\{a_i\}$ decay algebraically, Example 3 points out that the APE_{δ_n} is no longer asymptotically efficient if δ_n is bounded away from 1. To alleviate this difficulty, Theorem 2 (also in Section 3) allows δ_n to converge at a certain rate to 1 and offers a theoretical justification for the proposed modification. In light of this result, a class of APE_{δ_n} criteria that can achieve the asymptotic efficiency in both exponential- and algebraic-decay cases is given; see Examples 4 and 5 after Theorem 2.

The second issue concerns the performances of the information criterion and its relation to the APE_{δ_n} from the same-realization prediction point of view. The value of the information criterion for model $\text{AR}(k)$ is defined by

$$\text{IC}_{P_n}(k) = \log \hat{\sigma}_n^2(k) + \frac{P_n k}{n}, \quad (1.8)$$

where $P_n > 1$ is a positive number (possibly) depending on n , \log denotes the natural logarithm,

$$\hat{\sigma}_n^2(k) = \frac{1}{N} \sum_{t=K_n}^{n-1} (x_{t+1} + \hat{\mathbf{a}}'_n(k) \mathbf{x}_t(k))^2, \quad (1.9)$$

and $N = n - K_n$. Note that the AIC (Akaike, 1974), BIC (Schwarz, 1978) and HQ criterion (Hannan and Quinn, 1979) correspond to IC_{P_n} with $P_n = 2, \log n$, and $c \log_2 n$, respectively, where $c > 2$ and $\log_2 n = \log(\log n)$. Equation (1.8) is referred to as an AIC-like criterion if P_n is independent of n , and as a BIC-like criterion if $P_n \rightarrow \infty$ and $P_n = o(n)$. With the help of Proposition 2, Theorem 3 (see Section 4) gives an asymptotic expression for $q_n(\hat{k}_{n,P_n})$, where

$$\hat{k}_{n,P_n} = \arg \min_{1 \leq k \leq K_n} \text{IC}_{P_n}(k). \quad (1.10)$$

This result extends Corollary 1 of Ing and Wei (2005), which only focuses on the MSPE of the AIC-like criteria. An interesting implication of Theorem 3 is that the

HQ criterion is asymptotically efficient in the exponential-decay case whereas the BIC is not; see Examples 6 and 7 in Section 4. While both the HQ and BIC are known to be consistent in finite-order AR models (Hannan and Quinn, 1979), these examples show that their prediction performances can differ remarkably in the AR(∞) case. Based on Theorems 1-3, an asymptotic equivalence between IC_{P_n} and APE_{δ_n} , with δ_n and P_n satisfying (4.8), is given at the end of Section 4; see (4.9).

The third issue in which we are interested is a long-standing unresolved problem concerning time series model selection. Under the assumption that (1.1) does not degenerate to an AR model of finite order, Ing and Wei (2005) recently showed that AIC, satisfying (2.3), is asymptotically efficient for same-realization predictions. However, if the order of the underlying AR model is finite, then, as mentioned previously, the BIC-like criteria (e.g., HQ and BIC) are consistent, but AIC, which asymptotically will choose an overparameterized model with positive probability, does not possess this property (Shibata, 1976). When the APE_{δ_n} criteria are used instead, the choice between $\delta_n \rightarrow 0$ and $\delta_n \rightarrow 1$ also leads to the same difficulty; see Remark 5 in Section 3. To tackle these dilemmas, in Section 5, we first concentrate on an important special case where $\{a_i\}$ either decay exponentially or are zero for all but a finite number of i . It is shown in Theorem 5 that $IC_{P_n}(k)$, with $P_n \rightarrow \infty$ and $P_n = o(\log n)$ and $APE_{\delta_n}(k)$, with $\delta_n^{-1} \rightarrow \infty$ and $\log \delta_n^{-1} = o(\log n)$, can simultaneously achieve asymptotic efficiency over these two types of AR processes. However, if the case where $\{a_i\}$ decay algebraically is also included, then the criteria proposed by Theorem 5 fail to preserve the same optimality. A two-stage procedure, (5.1), which is a hybrid between AIC and a BIC-like criterion, is provided as a remedy. Its validity is justified theoretically in Theorem 6 (also in Section 5). Note that the results mentioned above are verified under the assumption that all moments of e_t are finite (see (K.3) in Section 2). It is made to obtain a uniform moment bound for $\hat{R}_i^{-1}(k)$ (see (B.6) in Appendix B), based on the recent work of Ing and Wei (2003, Theorem 2; 2005, Proposition 1). In fact, (K.3) can be slightly relaxed at the cost of reducing the number of candidate models. However, the details are not pursued here in order to simplify the discussion. Simulation results illustrating finite sample performances of the aforementioned criteria are given in Section 6. For ease of reading, the proofs of the results in Sections 2-5 are deferred to Appendices A-D, respectively.

2. PRELIMINARY RESULTS

We first list a set of assumptions that are used throughout the paper.

(K.1) Let $\{x_t\}$ be a linear process satisfying (1.1) with $A(z) = 1 + a_1z + a_2z^2 + \dots \neq 0$ for $|z| \leq 1$. Furthermore, let the coefficients $\{a_i\}$ obey $\sum_{i=1}^{\infty} |i^{1/2}a_i| < \infty$.

(K.2) Let the distribution function of e_t be denoted by F_t . There are two arbitrarily small positive numbers, α and δ_0^* , and one arbitrarily large positive number, C_0 , such that for all $t = \dots, -1, 0, 1, \dots$ and $|x - y| < \delta_0^*$,

$$|F_t(x) - F_t(y)| \leq C_0 |x - y|^\alpha.$$

(K.3) $\sup_{-\infty < t < \infty} E |e_t|^s < \infty$, $s = 1, 2, \dots$.

(K.4) The maximal order K_n satisfies

$$C_l \leq \frac{K_n^{2+\delta_1^*}}{n} \leq C_u,$$

where δ_1^* , C_l , and C_u are some prescribed positive numbers.

(K.5) $a_n \neq 0$ for infinitely many n .

First note that the MSPE of $\hat{x}_{n+1}(k)$, $q_n(k)$ (see (1.7)), can be expressed as

$$\sigma^2 + E (\mathbf{f}(k) + \mathcal{S}(k))^2, \quad (2.1)$$

where

$$\mathbf{f}(k) = \mathbf{x}'_n(k) \hat{R}_n^{-1}(k) \frac{1}{N} \sum_{j=K_n}^{n-1} \mathbf{x}_j(k) e_{j+1,k}, \quad e_{j+1,k} = x_{j+1} + \sum_{l=1}^k a_l(k) x_{j+1-l},$$

$$(a_1(k), \dots, a_k(k))' = \mathbf{a}(k) = \arg \min_{(c_1, \dots, c_k)' \in R^k} E \left(x_{k+1} + \sum_{l=1}^k c_l x_{k+1-l} \right)^2,$$

and

$$\mathcal{S}(k) = \sum_{i=1}^{\infty} (a_i - a_i(k)) x_{n+1-i}$$

with $a_i(k) = 0$ for $i > k$. To simplify the notation, $\mathbf{a}(k)$ is sometimes viewed as an infinite-dimensional vector with undefined entries set to zero. Ing and Wei (2003, Theorem 3) obtained an asymptotic expression for $q_n(k) - \sigma^2$, which holds uniformly for all $1 \leq k \leq K_n$. This result is summarized in the following proposition.

Proposition 1. *Assume that (K.1)–(K.4) hold. Then,*

$$\lim_{n \rightarrow \infty} \max_{1 \leq k \leq K_n} \left| \frac{q_n(k) - \sigma^2}{L_n(k)} - 1 \right| = 0, \quad (2.2)$$

where

$$L_n(k) = \frac{k\sigma^2}{N} + \|\mathbf{a} - \mathbf{a}(k)\|_R^2,$$

and for any infinite-dimensional vector $\mathbf{d} = (d_1, d_2, \dots)'$, $\|\mathbf{d}\|_R^2 = \sum_{i \leq j \leq \infty} d_i d_j \gamma_{i-j}$, with $\gamma_{i-j} = E(x_i x_j)$. Also note that $\|\mathbf{a} - \mathbf{a}(k)\|_R^2 = E(S^2(k))$.

The first term of $L_n(k)$, $k\sigma^2/N$, which is proportional to k , can be viewed as a measure of model complexity. The second term of $L_n(k)$, $\|\mathbf{a} - \mathbf{a}(k)\|_R^2$, which decreases as k increases, measures the goodness of fit. If one attempts to find an order k whose corresponding predictor, $\hat{x}_{n+1}(k)$, has the minimal MSPE, then some data-driven order selection criteria are needed. An order selection criterion, \hat{k}_n , is said to be asymptotically efficient if $\hat{x}_{n+1}(\hat{k}_n)$ satisfies

$$\limsup_{n \rightarrow \infty} \frac{q_n(\hat{k}_n) - \sigma^2}{\min_{1 \leq k \leq K_n} q_n(k) - \sigma^2} \leq 1, \quad (2.3)$$

where $1 \leq \hat{k}_n \leq K_n$. Inequality (2.3) says that the (second-order) MSPE of the predictor with order determined by an asymptotically efficient criterion is ultimately not greater than that of the best predictor among $\{\hat{x}_{n+1}(1), \dots, \hat{x}_{n+1}(K_n)\}$. In view of (2.2), (2.3) is equivalent to

$$\limsup_{n \rightarrow \infty} \frac{q_n(\hat{k}_n) - \sigma^2}{L_n(k_n^*)} \leq 1, \quad (2.4)$$

where $L_n(k_n^*) = \min_{1 \leq k \leq K_n} L_n(k)$.

Let $\text{OS}_n(k)$ be an order selection function, and

$$\hat{k}_{n,OS} = \arg \min_{1 \leq k \leq K_n} \text{OS}_n(k) \quad (2.5)$$

be the selected order. We shall provide sufficient conditions under which $q_n(\hat{k}_{n,OS}) - \sigma^2$ can be asymptotically expressed in terms of the $L_n(\cdot)$ function. Define

$$L_{n,D_n}(k) = \frac{(D_n - 1)k\sigma^2}{N} + \|\mathbf{a} - \mathbf{a}(k)\|_R^2, \quad (2.6)$$

where $D_n > 1$, and

$$k_{n,D_n}^* = \arg \min_{1 \leq k \leq K_n} L_{n,D_n}(k). \quad (2.7)$$

Proposition 2. *Assume that (K.1)–(K.4) hold. If there exists a sequence of positive numbers $\{D_n\}$, with $\liminf_{n \rightarrow \infty} D_n > 1$, such that*

$$\lim_{n \rightarrow \infty} (D_n - 1) \frac{E(\mathcal{S}(\hat{k}_{n,OS}) - \mathcal{S}(k_{n,D_n}^*))^2}{L_{n,D_n}(k_{n,D_n}^*)} = 0 \quad (2.8)$$

and

$$\lim_{n \rightarrow \infty} (D_n - 1) \frac{E(\mathbf{f}(\hat{k}_{n,OS}) - \mathbf{f}(k_{n,D_n}^*))^2}{L_{n,D_n}(k_{n,D_n}^*)} = 0, \quad (2.9)$$

where $\mathcal{S}(k)$ and $\mathbf{f}(k)$ are defined after (2.1), then

$$\lim_{n \rightarrow \infty} \frac{q_n(\hat{k}_{n,OS}) - \sigma^2}{L_n(k_{n,D_n}^*)} = 1. \quad (2.10)$$

Moreover, if

$$\lim_{n \rightarrow \infty} \frac{L_n(k_{n,D_n}^*)}{L_n(k_n^*)} = 1, \quad (2.11)$$

then (2.3) ((2.4)) holds for $\hat{k}_n = \hat{k}_{n,OS}$.

Remark 1. If (K.1), (K.5), $\sup_{-\infty < t < \infty} E|e_t|^4 < \infty$, and $K_n = o(n^{1/2})$ are assumed, and (2.8) and (2.9) are replaced with

$$\text{p-lim}_{n \rightarrow \infty} (D_n - 1) \frac{L_n(\hat{k}_{n,OS}) - L_n(k_{n,D_n}^*)}{L_{n,D_n}(k_{n,D_n}^*)} = 0, \quad (2.12)$$

then it is shown in Appendix A that

$$\text{p-lim}_{n \rightarrow \infty} \frac{E\{(y_{n+1} - \hat{y}_{n+1}(\hat{k}_{n,OS}))^2 | x_1, \dots, x_n\} - \sigma^2}{L_n(k_{n,D_n}^*)} = 1, \quad (2.13)$$

where y_{n+1} is the future value of $\{y_1, \dots, y_n\}$, which is a realization from an independent copy of $\{x_t\}$, and $\hat{y}_{n+1}(k) = -\mathbf{y}'_n(k)\hat{a}_n(k)$ with $\mathbf{y}'_n(k) = (y_n, \dots, y_{n+1-k})$. Note that (2.13) gives an asymptotic expression for the (conditional) MSPE of $\hat{k}_{n,OS}$ in independent-realization settings. For further discussion, see Remark 6 in Section 3.

□

Proposition 2 asserts that if $\hat{k}_{n,OS}$ is sufficiently close to k_{n,D_n}^* in the sense of (2.8) and (2.9), then $q_n(\hat{k}_{n,OS}) - \sigma^2$ has the asymptotic expression $L_n(k_{n,D_n}^*)$. In addition, if (2.11) also holds, then $\hat{k}_{n,OS}$ is asymptotically efficient. Proposition 2 plays a prominent role in justifying APE_{δ_n} 's and IC_{P_n} 's asymptotic (in)efficiency in various situations; see Sections 3-5. To apply Proposition 2, it is important to determine a penalty term D_n associated with the selection criterion $OS_n(k)$ and then justify (2.8) and (2.9) under suitable assumptions. For the first task, it is shown in Section 3 that the D_n associated with $APE_{\delta_n}(k)$ is

$$D_{APE_{\delta_n}} = 1 + \frac{N \log \delta_n^{-1}}{n(1 - \delta_n)}. \quad (2.14)$$

According to Appendix C, the D_n associated with $IC_{P_n}(k)$ and

$$\frac{S_n^{(P_n)}(k)}{N} = \left(1 + \frac{P_n}{N}\right) \hat{\sigma}_n^2(k) \quad (2.15)$$

is P_n . To facilitate the second task, inspired by Ing and Wei (2005, (3.9)), Assumption (K.6) (see below) is frequently used in the rest of this paper. As will be seen in Appendices B-D, it is introduced to deal with the complicated dependency conditions among the selected order, estimated parameters, and future observations. (Note that in the independent-realization settings, the future value to be predicted is independent of the selected order and estimated parameters.)

(K.6) For any $\xi > 0$, there is a nonnegative exponent $0 \leq \theta = \theta(\xi) < 1$ and a positive number $M = M(\xi)$ such that

$$\liminf_{n \rightarrow \infty} R_n(\xi, \theta, M) > 0, \quad (2.16)$$

where, with K_n obeying (K.4), D_n satisfying $\liminf D_n > 1$ and $D_n = o(n)$, and $A_{D_n, \theta, M} = \{k : 1 \leq k \leq K_n, |k - k_{n,D_n}^*| \geq M(k_{n,D_n}^*)^\theta\}$,

$$R_n(\xi, \theta, M) = \min_{k \in A_{D_n, \theta, M}} (k_{n,D_n}^*)^\xi \frac{N \{L_{n,D_n}(k) - L_{n,D_n}(k_{n,D_n}^*)\}}{(D_n - 1) |k - k_{n,D_n}^*|}.$$

If $\{x_t\}$ is an AR process of finite order (namely, (K.5) does not hold), then (2.16) automatically holds. On the other hand, if (K.5) holds instead, by arguments similar to those in Examples 1 and 2 and the Appendix of Ing and Wei (2005), it can also be shown that (2.16) is fulfilled in the following cases: (a) the exponential-decay case,

$$C_1 k^{-\theta_1} e^{-\beta k} \leq \|\mathbf{a} - \mathbf{a}(k)\|_R^2 \leq C_2 k^{\theta_1} e^{-\beta k}, \quad (2.17)$$

where $C_2 \geq C_1 > 0$, $\theta_1 \geq 0$, and $\beta > 0$ (note that if (K.1) is assumed, then (2.17) is equivalent to $C_1^* k^{-\theta_1} e^{-\beta k} \leq \sum_{i \geq k} a_i^2 \leq C_2^* k^{\theta_1} e^{-\beta k}$, for some $C_2^* \geq C_1^* > 0$); and (b) the algebraic-decay case,

$$(C_3 - M_1 k^{-\xi_1}) k^{-\beta} \leq \|\mathbf{a} - \mathbf{a}(k)\|_R^2 \leq (C_3 + M_1 k^{-\xi_1}) k^{-\beta}, \quad (2.18)$$

where $C_3, M_1 > 0$, $\xi_1 \geq 2$ and $\beta > 1 + \delta_1^*$ (recall that δ_1^* is defined in (K.4)). These facts reveal that (2.16) is quite reasonable from both practical and theoretical points of view, since it includes the ARMA model (which is the most used short-memory time series model by far) and the AR(∞) model with algebraically decaying coefficients (which is of much theoretical interest in the context of model selection) as special cases.

It is worth mentioning that when (K.1)-(K.6) are assumed, (2.8) and (2.9) were verified by Ing and Wei (2005, (5.75) and (5.74), respectively) in the special case where $\hat{k}_{n,OS} = \arg \min_{1 \leq k \leq K_n} S_n^{(2)}(k)$ and $D_n = 2$. Using similar arguments and assumptions, it can be shown that (2.8) and (2.9) are still valid for $\hat{k}_{n,OS} = \hat{k}_{n,P_n}$ (defined in (1.10)), $D_n = P_n$, and $1 < P_n = \alpha < \infty$ independent of n . However, to justify (2.8) and (2.9) in the case $\hat{k}_{n,OS} = \hat{k}_{n,\delta_n}$ and $D_n = D_{APE_{\delta_n}}$ or in the case $\hat{k}_{n,OS} = \hat{k}_{n,P_n}$ and $D_n = P_n \rightarrow \infty$, a much more delicate analysis is required. This problem is tackled in the next two sections. Note that in the finite-order AR models, (2.8) and (2.9) can also be verified for these two type of criteria under suitable assumptions; see Section 5 for more details.

As observed in Proposition 2, (2.11) is an important key to the asymptotic efficiency. It holds if the penalty term D_n satisfies

$$\lim_{n \rightarrow \infty} D_n = 2. \quad (2.19)$$

To see this, first observe that

$$\max_{1 \leq k \leq K_n} \left| \frac{L_{n,D_n}(k)}{L_n(k)} - 1 \right| \rightarrow 0, \quad (2.20)$$

if $D_n \rightarrow 2$. (2.20) and the fact that

$$1 \leq \frac{L_n(k_{n,D_n}^*)}{L_n(k_n^*)} = \frac{\frac{L_n(k_{n,D_n}^*)}{L_{n,D_n}(k_{n,D_n}^*)} L_{n,D_n}(k_{n,D_n}^*)}{\frac{L_n(k_n^*)}{L_{n,D_n}(k_n^*)} L_{n,D_n}(k_n^*)} \leq \frac{\frac{L_n(k_{n,D_n}^*)}{L_{n,D_n}(k_{n,D_n}^*)}}{\frac{L_n(k_n^*)}{L_{n,D_n}(k_n^*)}} \quad (2.21)$$

yield (2.11). When $\|\mathbf{a} - \mathbf{a}(k)\|_R^2$ decays exponentially or (K.5) is violated, (2.11) can hold without (2.19); see Example 1 in Section 3 and the proof of Theorem 4 in

Appendix D. For some other interesting discussions regarding (2.11), see Examples 2 and 3 in Section 3 and Examples 6-8 in Section 4.

3. The MSPE of APE_{δ_n} in $\text{AR}(\infty)$ processes

This section provides asymptotic expressions for $q_n(\hat{k}_{n,\delta_n}) - \sigma^2$. Without loss of generality, $n\delta_n, 1/n \leq \delta_n \leq 1 - (1/n)$, is assumed to be a positive integer. First note that

$$\text{APE}_{\delta_n}(k) = \sum_{i=n\delta_n}^{n-1} (x_{i+1} + \mathbf{x}'_i(k)\hat{\mathbf{a}}_i(k))^2 = \sum_{i=n\delta_n}^{n-1} \{e_{i+1} + \hat{e}_{i,k} + (e_{i+1,k} - e_{i+1})\}^2, \quad (3.1)$$

where $\hat{e}_{i,k} = \mathbf{x}'_i(k)(\hat{\mathbf{a}}_i(k) - \mathbf{a}(k))$ and $e_{i+1,k}$ is defined after (2.1). Following Lai and Wei (1982, (2.7)),

$$\begin{aligned} \sum_{i=n\delta_n}^{n-1} \hat{e}_{i,k}^2 &= \sum_{i=n\delta_n}^{n-1} h_i(k)e_{i+1,k}^2 + Q_{n\delta_n}(k) - Q_n(k) + \sum_{i=n\delta_n}^{n-1} h_i(k)\hat{e}_{i,k}^2 \\ &\quad - 2 \sum_{i=n\delta_n}^{n-1} (1 - h_i(k))\hat{e}_{i,k}e_{i+1,k}, \end{aligned} \quad (3.2)$$

where

$$h_i(k) = \mathbf{x}'_i(k) \left(\sum_{j=K_n}^i \mathbf{x}_j(k)\mathbf{x}'_j(k) \right)^{-1} \mathbf{x}_i(k)$$

and

$$Q_i(k) = \left(\sum_{j=K_n}^{i-1} \mathbf{x}_j(k)e_{j+1,k} \right)' \left(\sum_{j=K_n}^{i-1} \mathbf{x}_j(k)\mathbf{x}'_j(k) \right)^{-1} \left(\sum_{j=K_n}^{i-1} \mathbf{x}_j(k)e_{j+1,k} \right).$$

On substituting (3.2) into (3.1), one obtains

$$\begin{aligned} \text{APE}_{\delta_n}(k) &= \sum_{i=n\delta_n}^{n-1} e_{i+1}^2 + \left\{ \sum_{i=n\delta_n}^{n-1} h_i(k)e_{i+1,k}^2 - k\sigma^2 \log \delta_n^{-1} \right\} \\ &\quad + Q_{n\delta_n}(k) - Q_n(k) + \sum_{i=n\delta_n}^{n-1} h_i(k)\hat{e}_{i,k}^2 + 2 \sum_{i=n\delta_n}^{n-1} h_i(k)\hat{e}_{i,k}e_{i+1,k} \\ &\quad + \sum_{i=n\delta_n}^{n-1} \{(e_{i+1,k} - e_{i+1})^2 - \|\mathbf{a} - \mathbf{a}(k)\|_R^2\} + 2 \sum_{i=n\delta_n}^{n-1} (e_{i+1,k} - e_{i+1})e_{i+1} \\ &\quad + n(1 - \delta_n)L_{n,D_{\text{APE}_{\delta_n}}}(k), \end{aligned} \quad (3.3)$$

where $D_{\text{APE}_{\delta_n}}$ is defined in (2.14). When δ_n is bounded away from 1, Theorem 1 below provides sufficient conditions under which (2.8) and (2.9) hold for $\hat{k}_{n,OS} = \hat{k}_{n,\delta_n}$ and

$D_n = D_{APE_{\delta_n}}$. As a result, an asymptotic expression for $q_n(\hat{k}_{n,\delta_n}) - \sigma^2$ is obtained. Note that the relation $D_n = D_{APE_{\delta_n}}$ is used in the rest of this section.

Theorem 1. *Assume that (K.1)-(K.6) hold and $1/n \leq \delta_n \leq 1 - (1/n)$ satisfies*

$$\limsup_{n \rightarrow \infty} \delta_n < 1, \quad (3.4)$$

and

$$0 < \liminf_{n \rightarrow \infty} n^{\theta_3} \delta_n \leq \infty, \quad (3.5)$$

where $0 < \theta_3 < \delta_1^*/(2 + \delta_1^*)$. Moreover, if for some $0 \leq \theta = \theta(\xi) < 1$ and $\eta > 0$,

$$\lim_{n \rightarrow \infty} \frac{D_n - 1}{(k_{n,D_n}^*)^{1-\theta(1+\eta)}} = 0, \quad (3.6)$$

where θ is obtained from (K.6) when

$$0 < \xi < \min\{1/2, \{(2 + \delta_1^*)(1 - \theta_3)/2\} - 1\}, \quad (3.7)$$

then (2.8) and (2.9) are true for $\hat{k}_{n,OS} = \hat{k}_{n,\delta_n}$. Hence,

$$\lim_{n \rightarrow \infty} \frac{q_n(\hat{k}_{n,\delta_n}) - \sigma^2}{L_n(k_{n,D_n}^*)} = 1. \quad (3.8)$$

Remark 2. Note that θ in (3.6) is not uniquely determined. In order for (3.6) to be less stringent, θ can be chosen as small as possible; see Examples 1 and 3 below for more details. \square

Remark 3. Consider the following assumption:

(K.6') For any $\xi > 0$, there is a subsequence of $\{n\}$, $\{n_l\}$, such that

$$\liminf_{l \rightarrow \infty} R_{n_l}(\xi, 0, 1) > 0. \quad (3.9)$$

(Recall that $R_n(\xi, \theta, M)$ is defined in (K.6).) If, in place of (K.6), (K.6') is assumed in Theorem 1, then it can be shown that (3.8) remains valid for $n = n_l$, without imposing (3.6). This finding is applied in Example 2 below to illustrate that the APE_{δ_n} criteria, with δ_n decreasing to 0 at a polynomial rate, perform poorly in the case where the AR coefficients decay exponentially fast. \square

Remark 4. Under (K.5), it is not difficult to see that $k_{n,D_n}^* \rightarrow \infty$ as $n \rightarrow \infty$. Therefore, when $0 < \delta_n = \delta < 1$ is fixed with n , (3.6) automatically holds. \square

The following examples help gain a better understanding of Theorem 1.

Example 1. Assume that (K.1)-(K.4) hold and the AR coefficients satisfy

$$C_1 e^{-\beta k} \leq \|\mathbf{a} - \mathbf{a}(k)\|_R^2 \leq C_2 e^{-\beta k}, \quad (3.10)$$

where $0 < C_1 \leq C_2 < \infty$ and $\beta > 0$. Note that (3.10) is fulfilled by any causal and invertible ARMA(p, q) model with $q > 0$. As mentioned in Section 2, when (K.1) is assumed, (3.10) can also be expressed as $C'_1 e^{-\beta k} \leq \sum_{i \geq k} a_i^2 \leq C'_2 e^{-\beta k}$ for some $0 < C'_1 \leq C'_2 < \infty$. We shall show in this example that (3.8) follows if δ_n satisfies (3.4) and

$$\log \delta_n^{-1} = o(\log n). \quad (3.11)$$

Condition (3.11) guarantees (3.5). It can be shown to be equivalent to $n^\nu \delta_n \rightarrow \infty$ for all $\nu > 0$. Obviously, (3.4) and (3.11) are satisfied if $0 < \delta_n = \delta < 1$ is independent of n or if $\delta_n^{-1} = (\log n)^{\nu_1}$ for some $\nu_1 > 0$. Therefore, in view of Theorem 1 and the discussion given after (K.6), it remains to verify (3.6). By (3.10) and an argument similar to that used in the Appendix of Ing and Wei (2005), for some $C_1 > 0$,

$$\frac{1}{\beta} \log n - \frac{1}{\beta} \log(D_n - 1) - C_1 \leq k_{n, D_n}^* \leq \frac{1}{\beta} \log n - \frac{1}{\beta} \log(D_n - 1) + C_1, \quad (3.12)$$

and for any $\xi > 0$, (2.16) holds for $\theta = 0$ and some $M > 0$ (or for any $0 < \theta < 1$ and any $M > 0$). As a result, (3.6) holds for $\theta = 0$ and $\eta > 0$, and hence (3.8) follows. Moreover, (3.10) and the same argument used to prove (A.1) of Ing and Wei (2005) yield that for some $C_2 > 0$,

$$\frac{1}{\beta} \log n - C_2 \leq k_n^* \leq \frac{1}{\beta} \log n + C_2. \quad (3.13)$$

According to (3.10)-(3.13), we obtain (2.11), which together with (3.8), implies that the APE_{δ_n} , with δ_n satisfying (3.4) and (3.11), is asymptotically efficient. \square

Example 2. This example is given to indicate that if δ_n decays to 0 at a polynomial rate, then the APE_{δ_n} cannot be asymptotically efficient even in the exponential-decay case. More specifically, assume that (K.1)-(K.4) are satisfied,

$$\delta_n = C_1 n^{-\theta_3}, \quad (3.14)$$

where $C_1 > 0$ and $0 < \theta_3 < \delta_1^*/(2 + \delta_1^*)$, and the AR coefficients obey a special case of (3.10),

$$\|\mathbf{a} - \mathbf{a}(k)\|_R^2 = C_2 e^{-\beta k} (1 + G_k), \quad (3.15)$$

where $|G_k| < 1$, $G_k \rightarrow 0$ as $k \rightarrow \infty$, and C_2 and β are some positive numbers. By (3.15) and an argument similar to that used in Case II of Shibata (1980, p. 162), for sufficiently large n ,

$$k_{n,D_n}^* = m_{1,n} \text{ or } m_{1,n} + 1, \quad (3.16)$$

where $m_{1,n}$ is the largest integer $\leq m_{1,n}^* = (1/\beta) \log[C_2 N \beta / ((D_n - 1)\sigma^2)]$. Define $x_{1,n} = m_{1,n}^* - m_{1,n}$ and $z = (1/\beta) \log[\beta / (1 - e^{-\beta})]$. Since $0 < z < 1$, there is a positive number κ such that $0 < z - \kappa < z + \kappa < 1$. Define a set of positive integers $A_\kappa = \{n : |x_{1,n} - z| > \kappa, n = 1, 2, \dots\}$. Then, it can be shown that A_κ contains infinitely many elements. Moreover, for any $\xi > 0$ and any sequence of positive integers $\{n_l\} \subseteq A_\kappa$, (3.9) holds. Therefore, according to Remark 3, (3.8) is valid for $n = n_l$. By an analogy with (3.16), for sufficiently large n ,

$$k_n^* = m_{2,n} \text{ or } m_{2,n} + 1, \quad (3.17)$$

where $m_{2,n}$ is the largest integer $\leq (1/\beta) \log(C_2 N \beta / \sigma^2)$. (3.14)-(3.17) yield

$$\lim_{n \rightarrow \infty} \frac{L_n(k_{n,D_n}^*)}{L_n(k_n^*)} > 1, \quad (3.18)$$

which, together with (3.8) (with $n = n_l$), gives

$$\limsup_{n \rightarrow \infty} \frac{q_n(\hat{k}_{n,\delta_n}) - \sigma^2}{L_n(k_n^*)} > 1. \quad (3.19)$$

As a result, the APE_{δ_n} , with δ_n satisfying (3.14), fails to achieve (2.4) in the exponential-decay case. \square

Example 3. This example investigates the prediction performances of APE_{δ_n} in the algebraic-decay case (2.18). If (2.18), (3.4), and (3.5) are satisfied, then the same argument as the one in Example 2 of Ing and Wei (2005) yields that

$$k_{n,D_n}^* = \left(N C_3 \beta (D_n - 1)^{-1} \sigma^{-2} \right)^{1/(\beta+1)} + O(1), \quad (3.20)$$

and for any $\xi > 0$, (2.16) holds for any $1 - \min\{\xi, 1\} < \theta < 1$ and any $M > 0$. These facts and (3.5) guarantee that (3.6) is valid for $1 - \min\{\xi, 1\} < \theta < 1$ and $0 < \eta < (1 - \theta)/\theta$, where ξ satisfies (3.7). Consequently, when (K.1)-(K.4), (2.18), (3.4), and (3.5) are assumed, (3.8) is ensured by Theorem 1. By (A.9) of Ing and Wei (2005),

$$k_n^* = \left(N C_3 \beta \sigma^{-2} \right)^{1/(\beta+1)} + O(1). \quad (3.21)$$

This, (2.18), (3.4), (3.5), and (3.20) imply that

$$\liminf_{n \rightarrow \infty} \frac{L_n(k_{n,D_n}^*)}{L_n(k_n^*)} > 1, \quad (3.22)$$

According to (3.8) and (3.22), the APE_{δ_n} is not asymptotically efficient in this case.

□

As can be seen from Example 3, due to violation of (2.11), the APE_{δ_n} , with δ_n bounded away from 1, is not asymptotically efficient in the algebraic-decay case. In view of (2.19)-(2.21), a natural remedy for this difficulty is to let $\delta_n \rightarrow 1$. However, problems can still occur if δ_n converges "too fast" to 1. To see this, let $\delta_n = 1 - (1/n)$. Then, $\text{APE}_{\delta_n}(k) = (x_n - \hat{x}_n(k))^2$. Since models are determined only by the last period's prediction errors, it does not seem possible to establish any (asymptotically) optimal selection result in this case. To resolve this dilemma, some suitable choices of δ_n are introduced in Theorem 2. Some examples are also given after the theorem to help gain further insights into it.

Theorem 2. *Assume that (K.1)-(K.6) hold and $1/n \leq \delta_n \leq 1 - (1/n)$ satisfies $\lim_{n \rightarrow \infty} \delta_n = 1$. Moreover, if either of the following conditions holds, then (2.8) and (2.9) are valid for $\hat{k}_{n,OS} = \hat{k}_{n,\delta_n}$.*

- (i) $\lim_{n \rightarrow \infty} k_{n,D_n}^*/n^{\theta_3} = 0$ for any $\theta_3 > 0$ and $(1 - \delta_n)^{-1} = O(k_{n,D_n}^{*\xi_2})$ for some $0 < \xi_2 < 1/2$.
- (ii) $(1 - \delta_n)^{-1} = O(k_{n,D_n}^{*\xi_2})$ for some $0 < \xi_2 < \min\{1/2, \delta_1^*/2\}$.

Consequently, APE_{δ_n} is asymptotically efficient in the sense of (2.3) ((2.4)).

In light of Theorem 2, the following examples demonstrate how to choose δ_n such that the resulting APE_{δ_n} is asymptotically efficient in both the exponential- and algebraic-decay cases.

Example 4. Assume that (K.1)-(K.4) hold and the AR coefficients obey (2.17). Although Example 1 shows that when θ_1 in (2.17) is equal to 0, the APE_{δ_n} , with δ_n satisfying (3.4) and (3.11), is asymptotically efficient, it is unclear whether this result still holds for $\theta_1 > 0$. Fortunately, this difficulty can be bypassed by letting

$$\delta_n = 1 - C_1(\log n)^{-r}, \quad (3.23)$$

with $C_1 > 0$ and $0 < r < 1/2$. First note that under (2.17) and (3.23), the same argument as in Example 1 of Ing and Wei (2005) yields that for some $C_2 > 0$,

$$\frac{1}{\beta} \log n - C_2 \log_2 n \leq k_{n,D_n}^* \leq \frac{1}{\beta} \log n + C_2 \log_2 n, \quad (3.24)$$

and for any $\xi > 0$, (2.16) holds for any $0 < \theta < 1$ and any $M > 0$. Moreover, since condition (i) of Theorem 2 is ensured by (3.23) and (3.24), the APE_{δ_n} , with δ_n satisfying (3.23), is asymptotically efficient under (2.17). \square

Example 5. This example shows that if δ_n satisfies (3.23) with $C_1 > 0$ and $0 < r < \infty$, then the corresponding APE_{δ_n} is asymptotically efficient under the algebraic-decay case (2.18). To see this, first note that following the same line of reasoning as in Example 2 of Ing and Wei (2005), (3.20) is still valid, and for any $\xi > 0$, (2.16) holds for any $1 - \min\{\xi, 1\} < \theta < 1$ and any $M > 0$. In addition, since condition (ii) of Theorem 2 is ensured by (3.20) and the condition imposed on δ_n , the desired result follows from Theorem 2. \square

Examples 4 and 5 suggest that to achieve asymptotic efficiency through the APE_{δ_n} in both the exponential- and algebraic-decay cases, δ_n can be chosen to satisfy (3.23) with $C_1 > 0$ and $0 < r < 1/2$. However, the question of how to determine the best C_1 and r seems difficult to answer from a finite sample point of view. For some simulation results illustrating the APE_{δ_n} 's performances in finite samples, see Section 5. We close this section with two remarks concerning the performance of APE_{δ_n} in finite-order AR models and for independent-realization predictions.

Remark 5. When (1.1) degenerates to an $\text{AR}(p_0)$ model with $1 \leq p_0 < \infty$, it can be shown that \hat{k}_{n,δ_n} , with $\liminf_{n \rightarrow \infty} \delta_n > 0$, is not a consistent estimator of p_0 (e.g., Inoue and Kilian, 2006). On the other hand, if $\delta_n \rightarrow 0$ at a certain rate, then the corresponding APE_{δ_n} is consistent and asymptotically efficient (see Appendix D). Since these results and Theorem 2 offer totally different suggestions for choosing δ_n , it becomes very challenging to achieve asymptotic efficiency through the APE_{δ_n} when (1.1) is allowed to degenerate to a finite autoregression. In Section 5, some selection criteria to remedy this difficulty are proposed. \square

Remark 6. Note that the APE_{δ_n} described in Theorem 2 is also asymptotically efficient for independent-realization predictions. By Corollary B.1 (see Appendix B),

$$\text{p-lim}_{n \rightarrow \infty} \frac{L_{n,D_n}(\hat{k}_{n,\delta_n})}{L_{n,D_n}(k_{n,D_n}^*)} - 1 = 0. \quad (3.25)$$

Armed with (3.25) and (2.19)-(2.21), it can be shown that (2.12) holds for $\hat{k}_{n,OS} = \hat{k}_{n,\delta_n}$. Consequently, Remark 1 and (2.11) guarantee that

$$\text{p-lim}_{n \rightarrow \infty} \frac{E\{(y_{n+1} - \hat{y}_{n+1}(\hat{k}_{n,\delta_n}))^2 | x_1, \dots, x_n\} - \sigma^2}{L_n(k_n^*)} = 1, \quad (3.26)$$

which gives the claimed result. For more details on the definition of asymptotic efficiency in independent-realization settings, see Shibata (1980), Bhansali (1996), and Karagrigoriou (1997). \square

4. The MSPE of the IC_{P_n} in $\text{AR}(\infty)$ processes.

In this section, prediction performances of the information criterion, $\text{IC}_{P_n}(k)$, $P_n > 1$, are investigated. When P_n is independent of n , Ing and Wei (2005, Corollary 1) obtained an asymptotic expression for $q_n(\hat{k}_{n,P_n}) - \sigma^2$, where \hat{k}_{n,P_n} , defined in (1.10), is the minimizer of $\text{IC}_{P_n}(k)$, with $1 \leq k \leq K_n$ and K_n satisfying (K.4). Theorem 3 below extends Ing and Wei's result to the case where P_n is allowed to tend to ∞ with n . Note that the relation, $D_n = P_n$, is used throughout this section.

Theorem 3. *Let (K.1)-(K.6) hold and P_n satisfy*

$$\liminf_{n \rightarrow \infty} P_n > 1, \quad (4.1)$$

and

$$P_n = O(n^{\theta_3}), \quad (4.2)$$

for some $0 < \theta_3 < \delta_1^*/(4 + 2\delta_1^*)$. Moreover, if (3.6) holds with (3.7) replaced by

$$0 < \xi < \{\delta_1^*/(4 + 2\delta_1^*)\} - \theta_3, \quad (4.3)$$

then (2.8) and (2.9) are true for $\hat{k}_{n,OS} = \hat{k}_{n,P_n}$. Consequently,

$$\lim_{n \rightarrow \infty} \frac{q_n(\hat{k}_{n,P_n}) - \sigma^2}{L_n(k_{n,D_n}^*)} = 1. \quad (4.4)$$

Remark 7. If in Theorem 3, (K.6') holds instead of (K.6), then it can be shown that (4.4) is still valid for $n = n_l$ (note that n_l is defined in (K.6')). In this case, condition (3.6) is not required. This result can be applied to verify that BIC is not asymptotically efficient in the exponential-decay case; see Example 7 for more details. \square

Remark 8. Since (K.5) is assumed, (3.6) holds automatically if $P_n = O(1)$. \square

The following examples illustrate implications of Theorem 3. Special emphasis is placed on comparing the predictive capabilities of three well-known information criteria, AIC, HQ and BIC, in various situations.

Example 6. Assume that (K.1)-(K.4) hold and the AR coefficients satisfy (3.10). We shall show that $\text{IC}_{P_n}(k)$, with P_n satisfying (4.1) and

$$P_n = o(\log n), \quad (4.5)$$

is asymptotically efficient. Therefore, the AIC and HQ criterion are asymptotically efficient in this case. To see this, first note that the same reasoning as in Example 1 yields (3.12) and that for any $\xi > 0$, (2.16) holds for $\theta = 0$ and some $M > 0$ (or for any $0 < \theta < 1$ and any $M > 0$). These results and (4.5) imply that (3.6) holds for $\theta = 0$ and $\eta > 0$. According to Theorem 3, (4.4) follows. Consequently, the claimed result is ensured by (4.4) and by observing that (2.11) is valid under (3.10), (3.12), (3.13), and (4.5). \square

Example 7. This example illustrates that an information criterion cannot be asymptotically efficient in the exponential-decay case when the weight for penalizing the number of regressors in the model is "too strong". To see this, let (K.1)-(K.4) and (3.15) be satisfied and

$$P_n = C_1(\log n)^{C_2}, \quad (4.6)$$

where $C_1 > 0$ and $C_2 \geq 1$. Under these assumptions, (3.16) is obtained and (3.9) holds for any $\xi > 0$ and any sequence $\{n_l\} \subseteq A_\kappa$, where A_κ is defined as in Example 2. By Remark 7, (4.4) is valid for $n = n_l$. Moreover, since (3.15)-(3.17) and (4.6) yield (3.18), it is concluded that $\text{IC}_{P_n}(k)$, with P_n given by (4.6), is not asymptotically efficient. One important implication of this example is that BIC is not asymptotically efficient in the algebraic-decay case. \square

Example 8. Consider the algebraic-decay case, (2.18). Let P_n satisfy (4.1) and

$$P_n = O\left((\log n)^{C_1}\right), \quad (4.7)$$

for some $C_1 > 0$. By an argument similar to that used in Example 3, one obtains (3.20), and for any $\xi > 0$, (2.16) holds for any $1 - \min\{\xi, 1\} < \theta < 1$ and any $M > 0$. These facts and (4.7) imply that (3.6) is valid for $1 - \min\{\xi, 1\} < \theta < 1$ and $0 < \eta < (1 - \theta)/\theta$, where ξ satisfies (4.3). As a result, (4.4) follows from Theorem 3. Moreover, by (2.19)-(2.21), (2.11) holds when $\lim_{n \rightarrow \infty} P_n = 2$; and by (2.18), (3.20), and (3.21), $\limsup_{n \rightarrow \infty} L_n(k_{n,D_n}^*)/L_n(k_n^*) > 1$ when $\lim_{n \rightarrow \infty} P_n \neq 2$. This observation and (4.4) imply that AIC and AIC_C (Hurvich and Tsai, 1989) are asymptotically efficient in the algebraic-decay case, (2.18), whereas HQ, BIC and any information criterion with $\lim_{n \rightarrow \infty} P_n \neq 2$ are not. \square

As a final remark, note that when the conditions imposed by Theorems 1 and 3 (or Theorems 2 and 3) hold and

$$\lim_{n \rightarrow \infty} \frac{\log \delta_n^{-1}}{(1 - \delta_n)(P_n - 1)} = 1, \quad (4.8)$$

then

$$\lim_{n \rightarrow \infty} \frac{E \left(x_{n+1} - \hat{x}_{n+1}(\hat{k}_{n,P_n}) \right)^2 - \sigma^2}{E \left(x_{n+1} - \hat{x}_{n+1}(\hat{k}_{n,\delta_n}) \right)^2 - \sigma^2} = 1. \quad (4.9)$$

Instead of attempting to achieve a certain asymptotic optimality for prediction, (4.8) and (4.9) are interesting in that (4.8) can be used to connect the sequence δ_n defining the APE_{δ_n} with a corresponding parameter estimation penalty weight sequence, $P_n = 1 + (1 - \delta_n)^{-1} \log \delta_n^{-1}$, in such a way that $\hat{x}_{n+1}(\hat{k}_{n,P_n})$ and $\hat{x}_{n+1}(\hat{k}_{n,\delta_n})$ have the same asymptotic same-realization prediction (in)efficiency, as observed in (4.9). And, conversely, a sequence P_n implicitly determines a sequence δ_n through this same relation, which yields identical asymptotic (in)efficiency for the IC_{P_n} and APE_{δ_n} . This connection not only imparts to the APE_{δ_n} criteria the deep foundations of the information criteria, but also endows the information criteria with an on-line prediction meaning. For a related result, Wei (1992, Theorem 4.2.2), under (1.1) and certain moment conditions on e_t (which can be verified for the normal distribution), established an algebraic connection between the BIC and APE, $\log(\text{APE}(k)/n) = \text{BIC}(k) + o(\log n/n)$ a.s., where k is a positive integer and *fixed* with n . Therefore, except for the $o(\log n/n)$ term, the logarithm of $\text{APE}(k)/n$ is (a.s.) identical to $\text{BIC}(k)$. Hannan et al. (1989) also obtained the same result in a stationary $\text{AR}(p_0)$ model with $p_0 < \infty$ and $k \geq p_0$ (the correctly specified case). However, the equivalence introduced by (4.9) seems to be more relevant in situations where the two criteria's predictive capabilities after order selection are emphasized.

5. Optimal prediction for possibly degenerate $\text{AR}(\infty)$ processes

This section deals with optimal prediction problems in situations where the underlying $\text{AR}(\infty)$ process can degenerate to an AR process of finite order. We first adopt (K.5') to replace the truly infinite-order assumption, (K.5).

(K.5'): The AR coefficients satisfy either

- (i) $a_{p_0} \neq 0$ for some unknown $1 \leq p_0 < \infty$ and $a_l = 0$ for all $l \geq p_0 + 1$ or
- (ii) (3.10).

From a practical point of view, (K.5') is reasonably flexible because it contains any causal and invertible $\text{ARMA}(p, q)$ model, with $p + q \geq 1$, as a special case. Before tackling order selection problems under (K.5'), a preliminary result is needed, which shows that APE_{δ_n} and IC_{P_n} , with δ_n and P_n satisfying certain conditions, are asymptotically efficient in finite-order cases.

Theorem 4. Assume that (K.1)-(K.4) and (i) of (K.5') hold. Then, (2.8) and (2.9) hold for $(\hat{k}_{n,OS}, D_n) = (\hat{k}_{n,\delta_n}, D_{APE_{\delta_n}})$ and (\hat{k}_{n,P_n}, P_n) , where δ_n satisfies $\delta_n^{-1} \rightarrow \infty$ and (3.5), and P_n satisfies $P_n \rightarrow \infty$ and $P_n/n \rightarrow 0$. In addition, (2.3) ((2.4)) is satisfied by these criteria.

Remark 9. Since Theorem 4 adopts $\{\text{AR}(1), \dots, \text{AR}(K_n)\}$ as the set of candidate models, where $K_n \rightarrow \infty$ at a certain rate, the true model $\text{AR}(p_0)$ is included asymptotically. Zheng and Loh (1997) also took this approach. However, unlike Theorem 4, their main concern is with the consistency in order selection. \square

When (ii) of (K.5') holds, Example 6 points out that the IC_{P_n} , with $P_n = o(\log n)$, possesses asymptotic efficiency. On the other hand, if (i) of (K.5') is true, then Theorem 4 shows that the IC_{P_n} , with $P_n \rightarrow \infty$ and $P_n/n \rightarrow 0$, is asymptotically efficient under (K.1)-(K.4). These results taken together suggest that the IC_{P_n} , with $P_n \rightarrow \infty$ and $P_n = o(\log n)$, simultaneously achieve (2.3) over the two types of AR processes defined in (i) and (ii) of (K.5'). According to Example 1 and Theorem 4, the APE_{δ_n} , with $\delta_n^{-1} \rightarrow \infty$ and $\log \delta_n^{-1} = o(\log n)$, also has this property. These discussions are now summarized in the following theorem.

Theorem 5. Assume that (K.1)-(K.4) and (K.5') hold. Then, (2.3) ((2.4)) holds for $\hat{k}_n = \hat{k}_{n,\delta_n}$ and \hat{k}_{n,P_n} , where δ_n satisfies $\delta_n^{-1} \rightarrow \infty$ and $\log \delta_n^{-1} = o(\log n)$, and P_n satisfies $P_n \rightarrow \infty$ and $P_n = o(\log n)$.

As pointed out in Examples 3 and 8, the criteria given by Theorem 5 fail to preserve asymptotic efficiency when (2.18) is included in (K.5'). To overcome this difficulty, we propose using an alternative criterion that chooses order $\hat{k}_n^{(\iota)}$:

$$\hat{k}_n^{(\iota)} = \hat{k}_{n,2} I_{\{\hat{k}_{n,P_n} \neq \hat{k}_{n^\iota, P_{n^\iota}}\}} + \hat{k}_{n,P_n} I_{\{\hat{k}_{n,P_n} = \hat{k}_{n^\iota, P_{n^\iota}}\}}, \quad (5.1)$$

where $0 < \iota < 1$, $P_n \rightarrow \infty$, $\hat{k}_{n^\iota, P_{n^\iota}} = \arg \min_{1 \leq k \leq K_{n^\iota}} \text{IC}_{P_{n^\iota}}(k)$, and

$$\text{IC}_{P_{n^\iota}}(k) = \log \hat{\sigma}_{n^\iota}^2(k) + \frac{P_{n^\iota} k}{n^\iota},$$

with $\hat{\sigma}_{n^\iota}^2(k) = (1/N_\iota) \sum_{j=K_{n^\iota}}^{n^\iota-1} (x_{j+1} + \hat{\mathbf{a}}_{n^\iota}'(k) \mathbf{x}_j(k))^2$, $N_\iota = n^\iota - K_{n^\iota}$,

$$\hat{\mathbf{a}}_{n^\iota}(k) = -\hat{R}_{\iota, n^\iota}^{-1}(k) (1/N_\iota) \sum_{j=K_{n^\iota}}^{n^\iota-1} \mathbf{x}_j(k) x_{j+1},$$

and $\hat{R}_{\iota, n^\iota}(k) = (1/N_\iota) \sum_{j=K_{n^\iota}}^{n^\iota-1} \mathbf{x}_j(k) \mathbf{x}_j'(k)$ (note that without loss of generality, n^ι and K_{n^ι} are assumed to be positive integers). As observed, (5.1) is a hybrid selection procedure that combines together the AIC and a BIC-like criterion. If the true order

is finite, then it is expected that the orders selected by the BIC-like criterion at stages n^ι and n will ultimately be the same due to consistency. On the other hand, when the true order is infinite, an interesting result is derived for which it is nearly impossible for the BIC-like criterion to choose the same order at these different stages; see Appendix D. Therefore, it is reasonable to adopt $\hat{k}_{n,2}$ (the order selected by AIC) if the IC_{P_n} and $IC_{P_{n^\iota}}$ determine different orders, and \hat{k}_{n,P_n} (the order selected by the BIC-like criterion) otherwise. Theorem 6 justifies the validity of $\hat{k}_n^{(\iota)}$.

Theorem 6. *Let (K.1)-(K.4) and (K.6) (with $D_n = 2$) hold, and ι and P_n in (5.1) satisfy $0 < \iota < 1$, $P_n \rightarrow \infty$, $P_n = O(n^{\iota_1})$, with $0 < \iota_1 < \delta_1^*/(2 + \delta_1^*)$, and $P_n/P_{n^\nu} = O(1)$ for some $\nu > 0$. Further, assume that the AR coefficients meet either of the following conditions of*

(i) (i) of (K.5') or

(ii) for any $\xi > 0$, there is a nonnegative exponent, $0 \leq \theta = \theta(\xi) < 1$, and a positive number, $M = M(\xi) > 0$, such that

$$\liminf_{n \rightarrow \infty} \min_{k \in A_{P_n, \theta, M}} (k_{n, P_n}^*)^\xi \frac{L_{n, P_n}(k) - L_{n, P_n}(k_{n, P_n}^*)}{L_{n, P_n}(k_{n, P_n}^*)} > 0, \quad (5.2)$$

and for all sufficiently large n ,

$$A_{P_n^\iota, \theta, M}^C \cap A_{P_n, \theta, M}^C = \emptyset, \quad (5.3)$$

where $A_{P_n, \theta, M}$ is defined in (K.6), \emptyset denotes the empty set,

$$A_{P_n, \theta, M}^C = \{k : 1 \leq k \leq K_n, k \notin A_{P_n, \theta, M}\},$$

and

$$A_{P_{n^\iota}, \theta, M}^C = \{k : 1 \leq k \leq K_{n^\iota}, k \notin A_{P_{n^\iota}, \theta, M}\}.$$

(Note that (5.3) implicitly implies that $a_l \neq 0$ for infinitely many l .)

Then, (2.3) ((2.4)) holds for $\hat{k}_n = \hat{k}_n^{(\iota)}$.

As an application of Theorem 6, it is shown in Example 9 below that $\hat{k}_n^{(\iota)}$, $0 < \iota < 1$, is asymptotically efficient when the true model is either (i) an AR process of finite order, (ii) an AR(∞) process with coefficients satisfying (3.10) (the exponential-decay case), or (iii) an AR(∞) process with coefficients satisfying (2.18) (the algebraic-decay case). To simplify the discussion, let P_n be given by (4.6) with $C_1, C_2 > 0$, which satisfies all requirements for P_n imposed by Theorem 6.

Example 9. Assume that (K.1)-(K.4) hold, and either (K.5') or (2.18) is satisfied. To show that $\hat{k}_n^{(\iota)}$, $0 < \iota < 1$, is asymptotically efficient in this situation, in view of Theorem 6 and the discussion after (K.6), it suffices to show that (5.2) and (5.3) are guaranteed by (3.10) as well as (2.18). First, assume that (3.10) is true. Then, Example 6 shows that (3.12), with $D_n = P_n$, is valid, which further implies that for any $\xi > 0$, (5.2) holds for any $1 - \min\{\xi, 1\} < \theta < 1$ and any $M > 0$. In addition, (5.3) follows from (3.12) (with $D_n = P_n$), (4.6) (with $C_1, C_2 > 0$), $0 < \iota < 1$, and $0 < \theta < 1$. Next, let (2.18) hold. Reasoning as for Example 8, we obtain (3.20) (with $D_n = P_n$) and that for any $\xi > 0$, (5.2) holds for any $1 - \min\{\xi, 1\} < \theta < 1$ and any $M > 0$. Moreover, (5.3) is guaranteed by (3.20) (with $D_n = P_n$), (4.6) (with $C_1, C_2 > 0$), $0 < \iota < 1$, and $0 < \theta < 1$. Consequently, the desired result follows. \square

6. Simulation results.

To illustrate the practical implications of our theoretical results, a simulation study is conducted in this section. Let observations be generated from the ARMA(1,1) model

$$x_{t+1} = \phi_0 x_t + \varepsilon_t + \theta_0 \varepsilon_{t-1},$$

where ε_t 's are independent and identically $\mathcal{N}(0, 1)$ distributed and $(\phi_0, \theta_0) = (0.0, 0.98)$, $(0.5, 0.8)$, $(0.5, 0.4)$, and $(0.9, 0)$. For each combination of (ϕ_0, θ_0) , the empirical estimates of

$$RE(\hat{k}_n) = \frac{E \left(x_{n+1} - \hat{x}_{n+1}(\hat{k}_{n,2}) \right)^2 - 1}{E \left(x_{n+1} - \hat{x}_{n+1}(\hat{k}_n) \right)^2 - 1}$$

denoted by $\hat{RE}(\hat{k}_n)$ are obtained based on 5000 replications for $n = 180, 300, 500$, and 1000, where $\hat{k}_n = \hat{k}_{n,\delta_n}, \hat{k}_{n,P_n}, \hat{k}_n^{(\iota)}$, with $\delta_n = (\log n)^{-1}, 1 - (2/3)(\log n)^{-0.1}, 1 - (2/3)(\log n)^{-0.12}, 1 - (2/3)(\log n)^{-0.14}$, $P_n = 2.001 \log_2 n$ (HQ), $\log n$ (BIC), and $\iota = 0.69, 0.72, 0.75$. The penalty term of the BIC-like criterion associated with $\hat{k}_n^{(\iota)}$ is given by (4.6) with $C_1 = 0.8$ and $C_2 = 1$. In addition, K_n and K_{n^ι} are set to the largest integers less than or equal to $n^{1/2}$ and $n^{\iota/2}$, respectively. Obviously, $RE(\hat{k}_n)$ measures the relative prediction efficiency of $\hat{x}_{n+1}(\hat{k}_n)$ to $\hat{x}_{n+1}(\hat{k}_{n,2})$ (AIC), and $RE(\hat{k}_n) > 1$ ($RE(\hat{k}_n) < 1$) suggests that \hat{k}_n performs better (worse) than $\hat{k}_{n,2}$. These empirical results (see Table I) are summarized as follows:

(1) AIC and BIC: The relative efficiencies of AIC and BIC are clearly affected by the magnitude of the MA parameter in finite-sample situations. Table I shows that when $\theta_0 \geq 0.8$, AIC notably outperforms BIC, which coincides with our theoretical

findings in Examples 7 and 8 that BIC is not asymptotically efficient in truly $\text{AR}(\infty)$ models. In contrast, values of $\hat{RE}(\hat{k}_{n,\log n})$ are larger than 1 when $\theta_0 = 0.4$. However, since these values rapidly decrease from 1.26 to 1.08 as n grows from 180 to 1000, the theoretical result just mentioned does not seem to be seriously violated. On the other hand, when $\theta_0 = 0$, values of $\hat{RE}(\hat{k}_{n,\log n})$ are larger than 3.5, and do not exhibit any decreasing trend. This matches the fact that BIC is consistent and asymptotically efficient in finite-order AR models (see Section 5), whereas AIC is not.

(2) HQ and $\text{APE}_{\delta_{1,n}}$, where $\delta_{1,n} = (\log n)^{-1}$: First note that the prediction efficiencies of these two criteria seem quite close. They perform comparably to AIC when $\theta_0 = 0.8$, and much better than it when $\theta_0 \leq 0.4$. This phenomenon can be explained by the fact that HQ and $\text{APE}_{\delta_{1,n}}$ are asymptotically efficient in both the finite-order AR model and the $\text{AR}(\infty)$ model with AR coefficients decaying exponentially (see Theorem 5). Their efficiencies, however, are smaller than AIC in the case $\theta_0 = 0.98$. Since it is difficult to distinguish between an MA(1) process with a very large MA coefficient and an $\text{AR}(\infty)$ process with AR coefficients decaying algebraically in finite samples, Examples 3 and 8 (which show that HQ and $\text{APE}_{\delta_{1,n}}$ are not asymptotically efficient in the algebraic-decay case) may explain why HQ and $\text{APE}_{\delta_{1,n}}$ perform worse than AIC when θ_0 is very close to unity. In addition, we also observe that these two criteria are not as efficient as BIC in the case $\theta_0 = 0$, but they beat BIC in all other cases.

(3) $\text{APE}_{\delta_{i,n}}$, $i = 2, 3, 4$, where $\delta_{2,n} = 1 - (2/3)(\log n)^{-0.1}$, $\delta_{3,n} = 1 - (2/3)(\log n)^{-0.12}$, and $\delta_{4,n} = 1 - (2/3)(\log n)^{-0.14}$: Table I shows that $\text{APE}_{\delta_{i,n}}$, $i = 2, 3, 4$, holds a slight advantage (disadvantage) over AIC when $\theta_0 = 0.4$ ($\theta_0 \geq 0.8$). However, since the amount of the advantage (disadvantage) is not sizable, these Monte Carlo results seem to support the theoretical findings revealed in Examples 4-6 and 8 that AIC and these APE_{δ_n} criteria are asymptotically equivalent in both the exponential- and algebraic-decay cases. On the other hand, these criteria tend to outperform AIC when $\theta_0 = 0$. But they are still much less efficient than all other criteria in this case due to the lack of consistency in the finite-order case (see Remark 5).

(4) Two-stage criteria: One special feature of two-stage criteria is that they behave like AIC in situations where AIC dominates, and improve substantially over AIC in situations where AIC performs poorly. More specifically, values of $\hat{RE}(\hat{k}_n^{(l)})$ are rather close to 1 when $\theta_0 \geq 0.8$, and significantly larger than 1 when $\theta_0 = 0$. In the case $\theta_0 = 0.4$, the two-stage criteria perform slightly better than AIC when $n \leq 300$, and comparably to AIC when $n \geq 500$. These simulation results seem to match

quite well with the conclusion drawn from Example 9 that the two-stage criteria are asymptotically efficient in all three (finite-order, exponential-decay, and algebraic-decay) cases. When $\theta_0 = 0$, the prediction performances of the two-stage criteria are similar to those of HQ and $\text{APE}_{\delta_{1,n}}$ (particularly when n is large), but worse than those of BIC.

In conclusion, note that the finite-sample behaviors of the criteria considered in this section can be well predicted by the asymptotic results obtained in Sections 3-5. Some desirable features (when compared to AIC or BIC) of the $\text{APE}_{\delta_{i,n}}$, HQ, and two-stage criteria are particularly encouraging (see the discussions above). The tuning parameters adopted in this section may also serve as good initial values for pursuing better finite-sample efficiencies.

Appendix A: Proofs of Proposition 2 and (2.13).

PROOF OF PROPOSITION 2. In view of (1.7) and (2.1),

$$\frac{q_n(\hat{k}_{n,OS}) - \sigma^2}{L_n(k_{n,D_n}^*)}$$

TABLE I
EMPIRICAL ESTIMATES OF $RE(\hat{k}_n)$

n	Models (ϕ_0, θ_0)	APE_{δ_n}				IC		Two-stage		
		$\delta_{1,n}$	$\delta_{2,n}$	$\delta_{3,n}$	$\delta_{4,n}$	HQ	BIC	$n^{0.69}$	$n^{0.72}$	$n^{0.75}$
180	(0.0, 0.98)	0.88	0.93	0.92	0.93	0.89	0.78	0.95	0.94	0.94
	(0.5, 0.8)	0.95	0.95	0.95	0.94	0.98	0.83	0.98	0.97	0.97
	(0.5, 0.4)	1.28	1.07	1.05	1.03	1.36	1.26	1.08	1.08	1.08
	(0.9, 0.0)	2.21	1.34	1.33	1.28	2.31	3.59	1.81	1.86	1.95
300	(0.0, 0.98)	0.88	0.94	0.94	0.94	0.89	0.74	0.97	0.96	0.95
	(0.5, 0.8)	0.98	0.99	0.98	0.97	0.96	0.79	0.95	0.94	0.93
	(0.5, 0.4)	1.28	1.03	1.03	1.03	1.24	1.24	1.09	1.09	1.09
	(0.9, 0.0)	2.18	1.37	1.32	1.26	2.44	3.46	1.95	1.99	2.07
500	(0.0, 0.98)	0.85	0.94	0.95	0.95	0.85	0.68	0.96	0.95	0.94
	(0.5, 0.8)	0.97	0.97	0.97	0.96	0.98	0.78	0.97	0.95	0.95
	(0.5, 0.4)	1.28	1.10	1.05	1.04	1.32	1.17	1.03	1.02	1.06
	(0.9, 0.0)	2.31	1.36	1.31	1.27	2.64	4.17	2.39	2.43	2.41
1000	(0.0, 0.98)	0.86	0.95	0.96	0.95	0.86	0.66	0.99	0.98	0.98
	(0.5, 0.8)	1.05	0.97	0.96	0.96	1.01	0.80	0.97	0.97	0.95
	(0.5, 0.4)	1.36	1.12	1.09	1.04	1.37	1.08	1.00	1.00	0.98
	(0.9, 0.0)	2.33	1.27	1.26	1.21	2.86	4.07	2.65	2.74	2.67

Note: $\delta_{1,n} = (\log n)^{-1}$, $\delta_{2,n} = 1 - (2/3)(\log n)^{-0.1}$, $\delta_{3,n} = 1 - (2/3)(\log n)^{-0.12}$, and $\delta_{4,n} = 1 - (2/3)(\log n)^{-0.14}$.

$$= \frac{E\{\mathbf{f}(\hat{k}_{n,OS}) - \mathbf{f}(k_{n,D_n}^*) + \mathcal{S}(\hat{k}_{n,OS}) - \mathcal{S}(k_{n,D_n}^*) + \mathbf{f}(k_{n,D_n}^*) + \mathcal{S}(k_{n,D_n}^*)\}^2}{L_n(k_{n,D_n}^*)}. \quad (\text{A.1})$$

It is also not difficult to see that

$$\frac{L_{n,D_n}(k_{n,D_n}^*)}{L_n(k_{n,D_n}^*)} = O(D_n - 1). \quad (\text{A.2})$$

By (A.1), (A.2), (2.2), (2.8), and (2.9), (2.10) follows. Moreover, if (2.11) is assumed, (2.10) can be rewritten as

$$\lim_{n \rightarrow \infty} \frac{q_n(\hat{k}_{n,OS}) - \sigma^2}{L_n(k_n^*)} = 1, \quad (\text{A.3})$$

and hence (2.3) (or (2.4)) holds for $\hat{k}_n = \hat{k}_{n,OS}$. \square

PROOF OF (2.13). Under (K.1), (K.5), $\sup_{-\infty < t < \infty} E|e_t|^4 < \infty$, and $K_n = o(n^{1/2})$, an argument given in Ing and Wei (2005, p. 2448) yields

$$\text{p-lim}_{n \rightarrow \infty} \frac{E\{(y_{n+1} - \hat{y}_{n+1}(\hat{k}_{n,OS}))^2 | x_1, \dots, x_n\} - \sigma^2}{L_n(\hat{k}_{n,OS})} = 1. \quad (\text{A.4})$$

The desired result now follows from (2.12) and (A.4). \square

Appendix B: Proofs of Theorems 1 and 2.

In the rest of this paper, C is used to denote a generic positive constant independent of the sample size n and of any index with an upper (or lower) limit dependent on n . It also may have different values in different places. We start with a modification of Lemma 6 of Ing and Wei (2005).

Lemma B.1. *Assume that (K.1) holds with $\sum_{i=1}^{\infty} |i^{1/2} a_i| < \infty$ replaced by $\sum_{i=1}^{\infty} |a_i| < \infty$ and $\sup_{-\infty < t < \infty} E|e_t|^{2q} < \infty$ for some $q \geq 2$. Let $\{m_{i,n}\}$, $i=0, 1, 2$, be sequences of positive integers satisfying $m_{2,n} \geq m_{1,n} \geq m_{0,n}$ for all $n \geq 1$. Then, for all $1 \leq k, j \leq m_{0,n}$,*

$$E \left| S_{m_{1,n}, m_{2,n}}^2(k) - \sigma_k^2 - \left(S_{m_{1,n}, m_{2,n}}^2(j) - \sigma_j^2 \right) \right|^q \leq C m_n^{-q/2} \|\mathbf{a}(j) - \mathbf{a}(k)\|_R^q, \quad (\text{B.1})$$

where $m_n = m_{2,n} - m_{1,n} + 1$, $S_{m_{1,n}, m_{2,n}}^2(k) = (1/m_n) \sum_{t=m_{1,n}}^{m_{2,n}} e_{t+1,k}^2$, $\mathbf{a}(j)$ and $\mathbf{a}(k)$ in (B.1) are viewed as infinite-dimensional vectors with undefined entries set to zero, and $\sigma_k^2 = E(e_{1,k}^2)$. Also note that $\|\mathbf{a}(j) - \mathbf{a}(k)\|_R^2 = \|\mathbf{a} - \mathbf{a}(j)\|_R^2 - \|\mathbf{a} - \mathbf{a}(k)\|_R^2$.

The proof of (B.1) is similar to that of Ing and Wei (2005, Lemma 6), and hence is omitted. Let $n\delta_n$, with $1/n \leq \delta_n \leq 1 - (1/n)$, be a positive integer. According to

(3.3), for $k \neq k_{n,D_n}^*$,

$$\begin{aligned} P(\hat{k}_{n,\delta_n} = k) &\leq P\left(\frac{\text{APE}_{\delta_n}(k)}{U_{\delta_n}(k)} \leq \frac{\text{APE}_{\delta_n}(k_{n,D_n}^*)}{U_{\delta_n}(k)}\right) \\ &\leq \sum_{l=1}^{12} P\left(|N_l(k)| \geq \frac{1}{12} V_{n,D_n}(k)\right), \end{aligned} \quad (\text{B.2})$$

where $D_n = D_{\text{APE}_{\delta_n}}$ is used throughout this appendix, $U_{\delta_n}(k) = n(1 - \delta_n)L_{n,D_n}(k)$,

$$\begin{aligned} V_{n,D_n}(k) &= (L_{n,D_n}(k) - L_{n,D_n}(k_{n,D_n}^*))L_{n,D_n}^{-1}(k), \\ |N_1(k)| &= U_{\delta_n}^{-1}(k) \left| \sum_{i=n\delta_n}^{n-1} h_i(k) e_{i+1,k}^2 - k\sigma^2 \log \delta_n^{-1} \right|, \\ |N_2(k)| &= U_{\delta_n}^{-1}(k) \left| \sum_{i=n\delta_n}^{n-1} h_i(k_{n,D_n}^*) e_{i+1,k_{n,D_n}^*}^2 - k_{n,D_n}^* \sigma^2 \log \delta_n^{-1} \right|, \\ |N_3(k)| &= U_{\delta_n}^{-1}(k) |Q_{n\delta_n}(k) - k\sigma^2|, \quad |N_4(k)| = U_{\delta_n}^{-1}(k) |Q_{n\delta_n}(k_{n,D_n}^*) - k_{n,D_n}^* \sigma^2|, \\ |N_5(k)| &= U_{\delta_n}^{-1}(k) |Q_n(k) - k\sigma^2|, \quad |N_6(k)| = U_{\delta_n}^{-1}(k) |Q_n(k_{n,D_n}^*) - k_{n,D_n}^* \sigma^2|, \\ |N_7(k)| &= U_{\delta_n}^{-1}(k) \left| \sum_{i=n\delta_n}^{n-1} h_i(k) \hat{e}_{i,k}^2 \right|, \quad |N_8(k)| = U_{\delta_n}^{-1}(k) \left| \sum_{i=n\delta_n}^{n-1} h_i(k_{n,D_n}^*) \hat{e}_{i,k_{n,D_n}^*}^2 \right|, \\ |N_9(k)| &= 2U_{\delta_n}^{-1}(k) \left| \sum_{i=n\delta_n}^{n-1} h_i(k) \hat{e}_{i,k} e_{i+1,k} \right|, \\ |N_{10}(k)| &= 2U_{\delta_n}^{-1}(k) \left| \sum_{i=n\delta_n}^{n-1} h_i(k_{n,D_n}^*) \hat{e}_{i,k_{n,D_n}^*} e_{i+1,k_{n,D_n}^*} \right|, \\ |N_{11}(k)| &= U_{\delta_n}^{-1}(k) \left| \sum_{i=n\delta_n}^{n-1} \{ \varepsilon_{i+1}^2(k) - \varepsilon_{i+1}^2(k_{n,D_n}^*) - \|\mathbf{a} - \mathbf{a}(k)\|_R^2 + \|\mathbf{a} - \mathbf{a}(k_{n,D_n}^*)\|_R^2 \} \right|, \\ |N_{12}(k)| &= U_{\delta_n}^{-1}(k) \left| \sum_{i=n\delta_n}^{n-1} (e_{i+1,k} - e_{i+1,k_{n,D_n}^*}) e_{i+1} \right|, \end{aligned}$$

and $\varepsilon_{i+1}(k) = e_{i+1,k} - e_{i+1}$.

By (B.2), Chebyshev's inequality, and moment bounds for $|N_i|$, $i = 1, \dots, 12$ (see Lemmas B.2-B.4 below), an upper bound for $P(\hat{k}_{n,\delta_n} = k)$ can be obtained. This upper bound plays an important role in verifying Theorems 1 and 2.

Lemma B.2. *Let the assumptions of Proposition 1 hold and $1/n \leq \delta_n \leq 1 - (1/n)$ satisfy (3.5). Then, for $q > 0$, all $1 \leq k \leq K_n$, and all sufficiently large n ,*

$$E(|N_1(k)|^q) \leq CU_{\delta_n}^{-q}(k) \left\{ \frac{k^{2q}}{(n\delta_n)^{q/2}} + (\log \delta_n^{-1})^q k^q \|\mathbf{a} - \mathbf{a}(k)\|_R^{2q} \right\}, \quad (\text{B.3})$$

and

$$E(|N_2(k)|^q) \leq CU_{\delta_n}^{-q}(k) \left\{ \frac{k_{n,D_n}^{*2q}}{(n\delta_n)^{q/2}} + (\log \delta_n^{-1})^q k_{n,D_n}^{*q} \|\mathbf{a} - \mathbf{a}(k_{n,D_n}^*)\|_R^{2q} \right\}. \quad (\text{B.4})$$

PROOF. We only prove (B.3) because the proof of (B.4) is similar. Define $D(i, k) = \mathbf{x}'_i(k)R^{-1}(k)\mathbf{x}_i(k)(i+1-K_n)^{-1}$, and $E(i, k) = k(i+1-K_n)^{-1}$. Then,

$$\begin{aligned} U_{\delta_n}(k)|N_1(k)| &\leq \left| \sum_{i=n\delta_n}^{n-1} (h_i(k) - D(i, k))e_{i+1,k}^2 \right| \\ &+ \left| \sum_{i=n\delta_n}^{n-1} D(i, k)(e_{i+1,k}^2 - e_{i+1}^2) \right| + \left| \sum_{i=n\delta_n}^{n-1} D(i, k)(e_{i+1}^2 - \sigma^2) \right| \\ &+ \left| \sum_{i=n\delta_n}^{n-1} (D(i, k) - E(i, k))\sigma^2 \right| + \sigma^2 \left| \sum_{i=n\delta_n}^{n-1} E(i, k) - k \log \delta_n^{-1} \right| \\ &\equiv I(k) + II(k) + III(k) + IV(k) + V(k). \end{aligned} \quad (\text{B.5})$$

By (3.5) and Ing and Wei (2005, Proposition 1), we have, for any $q > 0$, all $n\delta_n \leq i \leq n-1$, all $1 \leq k \leq K_n$ and all sufficiently large n ,

$$E\|\hat{R}_{i+1}^{-1}(k) - R^{-1}(k)\|^q \leq C \frac{k^q}{(i+1-K_n)^{q/2}}. \quad (\text{B.6})$$

Using Wei (1987, Lemma 2) and Jensen's inequality, it follows that for any $r > 0$, all $n\delta_n \leq i \leq n-1$, and all $1 \leq k \leq K_n$,

$$E(\|\mathbf{x}_i(k)\|^r) \leq Ck^{r/2}, \quad (\text{B.7})$$

and

$$E|e_{i+1,k}|^r \leq C. \quad (\text{B.8})$$

According to (B.6)-(B.8), Minkowski's inequality, and Hölder's inequality, we have, for $q \geq 1$, all $1 \leq k \leq K_n$, and all sufficiently large n ,

$$E(I(k))^q \leq \left(\sum_{i=n\delta_n}^{n-1} \left\| (h_i(k) - D(i, k))e_{i+1,k}^2 \right\|_q \right)^q \leq Ck^{2q}(n\delta_n)^{-q/2}, \quad (\text{B.9})$$

where for random variable z and positive number s , $\|z\|_s = E(|z|^s)^{1/s}$.

To deal with $II(k)$, note that the first moment bound theorem of Findley and Wei (1993) and Jensen's inequality yield for any $r > 0$, all $K_n \leq i \leq n - 1$, and all $1 \leq k \leq K_n$,

$$E(|\mathbf{x}'_i(k)R^{-1}(k)\mathbf{x}_i(k) - k|^r) \leq Ck^{r/2}. \quad (\text{B.10})$$

Reasoning as for (B.8), we have, for any $r > 0$, all $K_n \leq i \leq n - 1$ and all $1 \leq k \leq K_n$,

$$E(|\varepsilon_{i+1}(k)|^r) \leq C\|\mathbf{a} - \mathbf{a}(k)\|_R^r. \quad (\text{B.11})$$

(B.10), (B.11), Wei (1987, Lemma 2), and an argument similar to that used for obtaining (B.9) together imply that for $q \geq 2$ and all $1 \leq k \leq K_n$,

$$\begin{aligned} E(II(k))^q &\leq C \left\{ E \left| \sum_{i=n\delta_n}^{n-1} D(i, k) \varepsilon_{i+1}^2(k) \right|^q + E \left| \sum_{i=n\delta_n}^{n-1} D(i, k) \varepsilon_{i+1}(k) e_{i+1} \right|^q \right\} \\ &\leq C \left\{ (\log \delta_n^{-1})^q k^q \|\mathbf{a} - \mathbf{a}(k)\|_R^{2q} + (n\delta_n)^{-q/2} k^q \|\mathbf{a} - \mathbf{a}(k)\|_R^q \right\}. \end{aligned} \quad (\text{B.12})$$

Similarly,

$$E(III(k))^q \leq CE \left(\sum_{i=n\delta_n}^{n-1} D^2(i, k) \right)^{q/2} \leq Ck^q (n\delta_n)^{-q/2} \quad (\text{B.13})$$

holds for $q \geq 2$ and all $1 \leq k \leq K_n$.

To deal with $IV(k)$, it can be shown by some algebraic manipulations that

$$IV(k) = \sigma^2 \left| \frac{T_{n-1}(k)}{N} - \frac{T_{n\delta_n-1}(k)}{n\delta_n - K_n} + \sum_{i=n\delta_n}^{n-1} \frac{T_{i-1}(k)}{(i - K_n)(i + 1 - K_n)} \right|,$$

where $T_i(k) = \sum_{j=K_n}^i \mathbf{x}'_j(k)R^{-1}(k)\mathbf{x}_j(k) - k$. By an argument similar to that given in the proof of Lemma 3 of Ing and Wei (2005) and Jensen's inequality, one has for any $q > 0$, all $n\delta_n - 1 \leq i \leq n - 1$, and all $1 \leq k \leq K_n$,

$$E \left| \frac{T_i(k)}{i + 1 - K_n} \right|^q \leq C \frac{k^{3q/2}}{(i + 1 - K_n)^{q/2}}.$$

This and the Minkowski inequality yield that for $q \geq 1$ and all $1 \leq k \leq K_n$,

$$E(IV(k))^q \leq Ck^{3q/2} (n\delta_n)^{-q/2}. \quad (\text{B.14})$$

In addition, it is straightforward to show that for all $1 \leq k \leq K_n$,

$$V(k) \leq C \left(\frac{1 - \delta_n}{\delta_n} \right) k \left(\frac{K_n}{n} \right). \quad (\text{B.15})$$

Consequently, (B.3) follows from (B.5), (B.9), (B.12)-(B.15), Jensen's inequality, and the fact that for any $r > 0$,

$$\lim_{k \rightarrow \infty} k^r \|\mathbf{a} - \mathbf{a}(k)\|_R^{2r} = 0, \quad (\text{B.16})$$

which is ensured by (K.1). \square

Lemma B.3. *Under the assumptions of Lemma B.2, for $q > 0$, all $1 \leq k \leq K_n$, and all sufficiently large n ,*

$$E(|N_3(k)|^q) \leq CU_{\delta_n}^{-q}(k) \left\{ k^{2q}(n\delta_n)^{-q/2} + k^{q/2} \right\}, \quad (\text{B.17})$$

$$E(|N_4(k)|^q) \leq CU_{\delta_n}^{-q}(k) \left\{ k_{n,D_n}^{*2q}(n\delta_n)^{-q/2} + k_{n,D_n}^{*q/2} \right\}, \quad (\text{B.18})$$

$$E(|N_5(k)|^q) \leq CU_{\delta_n}^{-q}(k) \left(k^{2q}n^{-q/2} + k^{q/2} \right), \quad (\text{B.19})$$

$$E(|N_6(k)|^q) \leq CU_{\delta_n}^{-q}(k) \left(k_{n,D_n}^{*2q}n^{-q/2} + k_{n,D_n}^{*q/2} \right), \quad (\text{B.20})$$

$$E(|N_7(k)|^q) \leq CU_{\delta_n}^{-q}(k) k^{3q}(n\delta_n)^{-q}, \quad (\text{B.21})$$

$$E(|N_8(k)|^q) \leq CU_{\delta_n}^{-q}(k) k_{n,D_n}^{*3q}(n\delta_n)^{-q}, \quad (\text{B.22})$$

$$E(|N_9(k)|^q) \leq CU_{\delta_n}^{-q}(k) k^{2q}(n\delta_n)^{-q/2}, \text{ and} \quad (\text{B.23})$$

$$E(|N_{10}(k)|^q) \leq CU_{\delta_n}^{-q}(k) k_{n,D_n}^{*2q}(n\delta_n)^{-q/2}. \quad (\text{B.24})$$

PROOF. See Ing (2005, Lemmas A.7 and A.8).

Lemma B.4. *Let the assumptions of Lemma B.1 hold and $1/n \leq \delta_n \leq 1 - (1/n)$. Then, for $q \geq 2$ and all $1 \leq k \leq K_n$, with $K_n \leq n\delta_n$,*

$$E(|N_i(k)|^q) \leq CU_{\delta_n}^{-q/2}(k) L_{n,D_n}^{-q/2}(k) \|\mathbf{a}(k) - \mathbf{a}(k_{n,\delta_n}^*)\|_R^q, \quad (\text{B.25})$$

where $i = 11$ and 12 .

PROOF. First note that

$$E|L_{n,D_n}(k)N_{11}(k)|^q \leq E \left| \frac{\sum_{i=n\delta_n}^{n-1} \{e_{i+1,k}^2 - e_{i+1,k_{n,D_n}^*}^2 - \sigma_k^2 + \sigma_{k_{n,D_n}^*}^2\}}{n(1-\delta_n)} \right|^q$$

$$+ E \left| \frac{\sum_{i=n\delta_n}^{n-1} (e_{i+1,k} - e_{i+1,k_n^*,D_n}) e_{i+1}}{n(1-\delta_n)} \right|^q \equiv (I) + (II). \quad (\text{B.26})$$

According to (B.1), one has for all $1 \leq k \leq K_n$,

$$(I) \leq \frac{C \|\mathbf{a}(k) - \mathbf{a}(k_n^*,D_n)\|_R^q}{(1-\delta_n)^{q/2} n^{q/2}}. \quad (\text{B.27})$$

Lemma 2 of Wei (1987) and the convexity of $x^{q/2}, x > 0$, yield for all $1 \leq k \leq K_n$,

$$\begin{aligned} (II) &\leq \frac{C}{\{n(1-\delta_n)\}^{(q/2)+1}} \sum_{i=n\delta_n}^{n-1} E(|e_{i+1,k} - e_{i+1,k_n^*,D_n}|^q) \\ &\leq \frac{C \|\mathbf{a}(k) - \mathbf{a}(k_n^*,D_n)\|_R^q}{(1-\delta_n)^{q/2} n^{q/2}}. \end{aligned} \quad (\text{B.28})$$

Consequently, (B.25), with $i = 11$, is ensured by (B.26)-(B.28). The proof is completed by noting that (B.25), with $i = 12$, is an immediate consequence of (B.28). \square

Armed with Lemmas B.2-B.5, we have the following result.

Corollary B.1. *Let (K.1)-(K.5), (3.4), and (3.5) hold. Then, for any $r > 0$,*

$$\lim_{n \rightarrow \infty} E \left(\frac{L_{n,D_n}(\hat{k}_{n,\delta_n})}{L_{n,D_n}(k_n^*,D_n)} - 1 \right)^r = 0. \quad (\text{B.29})$$

PROOF. Define $I_{1,n}(k) = L_{n,D_n}(k)/L_{n,D_n}(k_n^*,D_n)$. Let $\epsilon > 0$ be arbitrarily given. Then, by (B.2), one has

$$\begin{aligned} E(I_{1,n}(\hat{k}_{n,\delta_n}) - 1)^r &= \sum_{k=1}^{K_n} (I_{1,n}(k) - 1)^r P(\hat{k}_{n,\delta_n} = k) \\ &\leq \epsilon^r + \sum_{l=1}^{12} \left\{ \sum_{k \in A_{\epsilon,n}^{(\delta_n)}} (I_{1,n}(k) - 1)^r P(|N_l(k)| \geq (1/12)V_{n,D_n}(k)) \right\}, \end{aligned} \quad (\text{B.30})$$

where $A_{\epsilon,n}^{(\delta_n)} = \{k : 1 \leq k \leq K_n, I_{1,n}(k) - 1 > \epsilon\}$. In view of (B.30), (B.29) holds if for $l = 1, \dots, 12$,

$$\lim_{n \rightarrow \infty} \sum_{k \in A_{\epsilon,n}^{(\delta_n)}} (I_{1,n}(k) - 1)^r P(|N_l(k)| > (1/12)V_{n,D_n}(k)) = 0. \quad (\text{B.31})$$

In the following, we only prove (B.31) for $l = 1, 3$, and 11 because the proofs for $l = 2, 7, 8, 9$, and 10 are similar to that for $l = 1$, proofs for $l = 4, 5$, and 6 are similar to that for $l = 3$, and the proof for $l = 12$ is similar to that for $l = 11$.

By (B.3), Chebyshev's inequality, (3.4), (3.5), and the facts that

$$L_{n,D_n}(k) \geq \|\mathbf{a} - \mathbf{a}(k)\|_R^2, \quad nL_{n,D_n}(k) \geq C \frac{k \log \delta_n^{-1}}{1 - \delta_n}, \quad (\text{B.32})$$

and $I_{1,n}(k) \leq C/L_{n,D_n}(k_{n,D_n}^*)$ if $1 \leq k \leq k_{n,D_n}^*$ and $I_{1,n}(k) \leq Ck/k_{n,D_n}^*$ if $k_{n,D_n}^* < k \leq K_n$, we have, for sufficiently large q ,

$$\begin{aligned} & \sum_{k \in A_{\epsilon,n}^{(\delta_n)}} (I_{1,n}(k) - 1)^r P(|N_1(k)| > (1/12)V_{n,D_n}(k)) \\ & \leq C \sum_{k \in A_{\epsilon,n}^{(\delta_n)}} I_{1,n}^r(k) V_{n,D_n}^{-(q-r)}(k) \{k^q f_{1,n}^{-q} + (f_{2,n}k)^q n^{-q}\} \\ & \leq C \left(\frac{1+\epsilon}{\epsilon}\right)^{q-r} \left[\sum_{k=1}^{k_{n,D_n}^*} \left\{ \frac{k^q (1-\delta_n)^r}{f_{1,n}^q (k_{n,D_n}^* f_{3,n} \delta_n)^r} + \frac{f_{2,n}^{q-r} k^q}{k_{n,D_n}^{*r} n^{q-r}} \right\} \right. \\ & \quad \left. + \sum_{k=k_{n,D_n}^*+1}^{K_n} \left\{ \frac{k^{q+r}}{k_{n,D_n}^{*r} f_{1,n}^q} + \frac{f_{2,n}^q k^{q+r}}{k_{n,D_n}^{*r} n^q} \right\} \right] = o(1), \quad (\text{B.33}) \end{aligned}$$

where $f_{1,n} = (\log \delta_n^{-1})(n\delta_n)^{1/2}$, $f_{2,n} = \log \delta_n^{-1}/(1 - \delta_n)$, and $f_{3,n} = \log \delta_n^{-1}/(n\delta_n)$. Therefore, (B.31) holds for $l = 1$.

By (B.17), (B.25), an argument similar to that used for obtaining (B.33) and the fact that $k_{n,D_n}^* \rightarrow \infty$ as $n \rightarrow \infty$, for sufficiently large q ,

$$\begin{aligned} & \sum_{k \in A_{\epsilon,n}^{(\delta_n)}} (I_{1,n}(k) - 1)^r P(|N_3(k)| > (1/12)V_{n,D_n}(k)) \\ & \leq C \left(\frac{1+\epsilon}{\epsilon}\right)^{q-r} \left\{ \sum_{k=1}^{K_n} I_{1,n}^r(k) k^q f_{1,n}^{-q} + \sum_{k=1}^{k_{n,D_n}^*} k^{q/2} U_{\delta_n}^{-q}(k_{n,D_n}^*) \right. \\ & \quad \left. + \sum_{k=k_{n,D_n}^*+1}^{K_n} k^{q/2} U_{\delta_n}^{-q}(k) I_{1,n}^r(k) \right\} = o(1), \quad (\text{B.34}) \end{aligned}$$

and

$$\begin{aligned} & \sum_{k \in A_{\epsilon,n}^{(\delta_n)}} (I_{1,n}(k) - 1)^r P(|N_{11}(k)| > (1/12)V_{n,D_n}(k)) \\ & \leq C \left(\frac{1+\epsilon}{\epsilon}\right)^{q-r} \left\{ \sum_{k=1}^{k_{n,D_n}^*} U_{\delta_n}^{-q/2}(k_{n,D_n}^*) + \sum_{k=k_{n,D_n}^*+1}^{K_n} U_{\delta_n}^{-q/2}(k) I_{1,n}^r(k) \right\} \\ & = o(1). \quad (\text{B.35}) \end{aligned}$$

In view of (B.33)-(B.35), the proof is complete. \square

Corollary B.2. *Assume that (K.1)-(K.6) hold and δ_n satisfies (3.4) and (3.5). Then, for sufficiently large q ,*

$$\begin{aligned} E \left| \frac{\mathcal{S}(\hat{k}_{n,\delta_n}) - \mathcal{S}(k_{n,D_n}^*)}{(L_{n,D_n}(\hat{k}_{n,\delta_n}))^{1/2}} \right|^{2q} &= O((k_{n,D_n}^*)^{-(1-\theta)q+\theta}) + o((\log \delta_n^{-1})^{-q}) \\ &+ O((\log \delta_n^{-1})^{-q/2}(k_{n,D_n}^*)^{(-q/2)+\theta}), \end{aligned} \quad (\text{B.36})$$

where $\mathcal{S}(k)$ is defined in Section 2 and $0 \leq \theta = \theta(\xi) < 1$ is obtained from (K.6) with ξ satisfying (3.7).

PROOF. Let ξ satisfy (3.7). Then, (K.6) guarantees that there are $0 \leq \theta = \theta(\xi) < 1$ and $M = M(\xi) > 0$ such that (2.16) is fulfilled. Define $I_{2,n}(k) = L_{n,D_n}(k) - L_{n,D_n}(k_{n,D_n}^*)$. By Hölder's inequality and the fact that for any $h > 0$,

$$\begin{aligned} E \left| \mathcal{S}(k) - \mathcal{S}(k_{n,D_n}^*) \right|^{2h} &\leq C \left\| \mathbf{a}(k) - \mathbf{a}(k_{n,D_n}^*) \right\|_R^{2h} \\ &\leq C \left(I_{2,n}(k) + f_{2,n} N^{-1} |k - k_{n,D_n}^*|^2 \right)^h \end{aligned} \quad (\text{B.37})$$

(which follows from Wei (1987, Lemma 2), (K.3), and the definition of k_{n,D_n}^*), one has for $q > 0$ and $1 < r < \infty$,

$$\begin{aligned} E \left| \frac{\mathcal{S}(\hat{k}_{n,\delta_n}) - \mathcal{S}(k_{n,D_n}^*)}{L_{n,D_n}^{1/2}(\hat{k}_{n,\delta_n})} \right|^{2q} &\leq \sum_{k=1}^{K_n} \left(E \left| \frac{\mathcal{S}(k) - \mathcal{S}(k_{n,D_n}^*)}{L_{n,D_n}^{1/2}(k)} \right|^{2qr} \right)^{\frac{1}{r}} P^{\frac{r-1}{r}}(\hat{k}_{n,\delta_n} = k) \\ &\leq C \sum_{k=1}^{K_n} \left\{ V_{n,D_n}^q(k) + \left| \frac{f_{2,n}(k - k_{n,D_n}^*)}{NL_{n,D_n}(k)} \right|^q \right\} P^{\frac{r-1}{r}}(\hat{k}_{n,\delta_n} = k) \\ &\leq C \left\{ \sum_{k=1}^{K_n} V_{n,D_n}^q(k) P^{\frac{r-1}{r}}(\hat{k}_{n,\delta_n} = k) + \sum_{\substack{k=1 \\ k \notin A_{D_n,\theta,M}}}^{K_n} \left| \frac{f_{2,n}(k - k_{n,D_n}^*)}{NL_{n,D_n}(k)} \right|^q \right. \\ &\quad \left. + \sum_{\substack{k=1 \\ k \in A_{D_n,\theta,M}}}^{K_n} \left| \frac{f_{2,n}(k - k_{n,D_n}^*)}{NL_{n,D_n}(k)} \right|^q P^{\frac{r-1}{r}}(\hat{k}_{n,\delta_n} = k) \right\} \equiv C\{(I) + (II) + (III)\}, \end{aligned} \quad (\text{B.38})$$

where $A_{D_n,\theta,M}$ is a set of positive integers defined in (K.6).

By the definitions of $A_{D_n,\theta,M}$, $L_{n,D_n}(k)$ and $L_{n,D_n}(k_{n,D_n}^*)$, it is easy to see that

$$(II) \leq C(k_{n,D_n}^*)^{-(1-\theta)q+\theta}. \quad (\text{B.39})$$

In view of (B.2) and the fact that for $a, b \geq 0$, $(a + b)^{(r-1)/r} \leq a^{(r-1)/r} + b^{(r-1)/r}$, one obtains

$$(I) \leq \sum_{l=1}^{12} \left\{ \sum_{k=1}^{K_n} V_{n,D_n}^q(k) P^{\frac{r-1}{r}} (|N_l(k)| \geq (1/12)V_{n,D_n}(k)) \right\}. \quad (\text{B.40})$$

In the following, we shall show that when q is sufficiently large,

$$\sum_{k=1}^{K_n} V_{n,D_n}^q(k) P^{\frac{r-1}{r}} (|N_l(k)| \geq (1/12)V_{n,D_n}(k)) = o((\log \delta_n^{-1})^{-q}), \quad (\text{B.41})$$

for $l = 1, \dots, 10$; and

$$\begin{aligned} & \sum_{k=1}^{K_n} V_{n,D_n}^q(k) P^{\frac{r-1}{r}} (|N_l(k)| \geq (1/12)V_{n,D_n}(k)) \\ &= O((\log \delta_n^{-1})^{-q/2} (k_{n,D_n}^*)^{(-q/2)+\theta}) + o((\log \delta_n^{-1})^{-q}), \end{aligned} \quad (\text{B.42})$$

for $l = 11$ and 12 . As a result, one has for sufficiently large q ,

$$(I) = O((\log \delta_n^{-1})^{-q/2} (k_{n,D_n}^*)^{(-q/2)+\theta}) + o((\log \delta_n^{-1})^{-q}). \quad (\text{B.43})$$

By Lemma B.2, (3.4), (3.5), (K.4), and (B.32), for sufficiently large q ,

$$\begin{aligned} & \sum_{k=1}^{K_n} V_{n,D_n}^q(k) P^{\frac{r-1}{r}} (|N_1(k)| \geq (1/12)V_{n,D_n}(k)) \leq C \sum_{k=1}^{K_n} [E\{|N_1(k)|^{\frac{qr}{r-1}}\}]^{\frac{r-1}{r}} \\ & \leq \frac{C}{(\log \delta_n^{-1})^q} \left(\sum_{k=1}^{K_n} \frac{k^q}{(n\delta_n)^{q/2}} + \frac{(\log \delta_n^{-1})^{2q} k^q}{(1 - \delta_n)^{qn^q}} \right) = o((\log \delta_n^{-1})^{-q}), \end{aligned} \quad (\text{B.44})$$

which yields (B.41) for $l = 1$. For $l = 3$, according to (3.5), Lemma B.3, (B.32), and the fact that $k_{n,D_n}^* \rightarrow \infty$ as $n \rightarrow \infty$, one has for sufficiently large q ,

$$\begin{aligned} & \sum_{k=1}^{K_n} V_{n,D_n}^q(k) P^{\frac{r-1}{r}} (|N_3(k)| \geq (1/12)V_{n,D_n}(k)) \leq C \sum_{k=1}^{K_n} [E\{|N_3(k)|^{\frac{qr}{r-1}}\}]^{\frac{r-1}{r}} \\ & \leq \frac{C}{(\log \delta_n^{-1})^q} \left(\sum_{k=1}^{K_n} \frac{k^q}{(n\delta_n)^{q/2}} + \sum_{k=1}^{k_{n,D_n}^*} \frac{k^{q/2}}{k_{n,D_n}^{*q}} + \sum_{k=k_{n,D_n}^*+1}^{K_n} k^{-q/2} \right) = o((\log \delta_n^{-1})^{-q}). \end{aligned} \quad (\text{B.45})$$

The proofs of (B.41) for $l = 2, 7, 8, 9$, and 10 are similar to that of (B.44) and the proofs of (B.41) for $l = 4, 5$ and 6 are similar to that of (B.45). We skip the details

in order to save space. The proof of (B.42) is a bit more complicated. By (2.16), Lemma B.4, (B.37), (3.4), and the restriction on ξ , one has for sufficiently large q ,

$$\begin{aligned}
& \sum_{k=1}^{K_n} V_{n,D_n}^q(k) P^{\frac{r-1}{r}} (|N_l(k)| \geq (1/12)V_{n,D_n}(k)) \\
\leq & \sum_{\substack{k=1 \\ k \notin A_{D_n,\theta,M}}}^{K_n} \{E|N_{11}(k)|^{\frac{qr}{r-1}}\}^{\frac{r-1}{r}} + \sum_{k \in A_{D_n,\theta,M}}^{K_n} V_{n,D_n}^{-q}(k) \{E|N_{11}(k)|^{\frac{2qr}{r-1}}\}^{\frac{r-1}{r}} \\
\leq & C \left\{ \sum_{\substack{k=1 \\ k \notin A_{D_n,\theta,M}}}^{K_n} \frac{\|\mathbf{a}(k) - \mathbf{a}(k_{n,D_n}^*)\|_R^q}{U_{\delta_n}^{q/2}(k) L_{n,D_n}^{q/2}(k)} + \sum_{k \in A_{D_n,\theta,M}}^{K_n} \frac{I_{2,n}^q(k) + |\frac{f_{2,n}(k - k_{n,D_n}^*)}{N}|^q}{U_{\delta_n}^q(k) I_{2,n}^q(k)} \right\} \\
\leq & C \left[\frac{k_{n,D_n}^{*\theta}}{(\log \delta_n^{-1} k_{n,D_n}^*)^{q/2}} + \frac{\{1 + (k_{n,D_n}^*)^{\xi q}\}}{(\log \delta_n^{-1})^q} \left\{ \sum_{k=1}^{k_{n,D_n}^*} k_{n,D_n}^{*-q} + \sum_{k=k_{n,D_n}^*+1}^{K_n} k^{-q} \right\} \right] \\
= & O((\log \delta_n^{-1})^{-q/2} (k_{n,D_n}^*)^{(-q/2)+\theta}) + o((\log \delta_n^{-1})^{-q}), \tag{B.46}
\end{aligned}$$

where $l = 11$ or 12 .

Following arguments similar to those used to obtain (B.40) and (B.44)-(B.46), it can be shown that

$$\begin{aligned}
(III) & \leq \sum_{l=1}^{12} \left\{ \sum_{\substack{k=1 \\ k \in A_{D_n,\theta,M}}}^{K_n} \left| \frac{f_{2,n}(k - k_{n,D_n}^*)}{(N L_{n,D_n}(k))} \right|^q P^{\frac{r-1}{r}} (|N_l(k)| \geq (1/12)V_{n,D_n}(k)) \right\} \\
& = o((\log \delta_n^{-1})^{-q}), \tag{B.47}
\end{aligned}$$

where the equality holds for sufficiently large q . (For a detailed proof of (B.47), see Ing (2005, Corollary A.2).) Consequently, (B.36) is ensured by (B.38), (B.39), (B.43), and (B.47). \square

Corollary B.3. *Assume that the assumptions of Corollary B.2 hold. Then, for sufficiently large q ,*

$$\lim_{n \rightarrow \infty} E \left| \frac{\mathbf{f}(\hat{k}_{n,\delta_n}) - \mathbf{f}(k_{n,D_n}^*)}{(L_{n,D_n}(\hat{k}_{n,\delta_n}))^{1/2}} \right|^{2q} = o((\log \delta_n^{-1})^q). \tag{B.48}$$

PROOF. (B.48) can be verified using arguments similar to those in the proofs of Ing and Wei (2005, Lemmas 7 and 8) and Corollary B.2 above. For details, see Ing (2005, Corollary A.3). \square

We are now ready to prove Theorem 1.

PROOF OF THEOREM 1. Let $q > \max\{\eta^{-1}, 1\}$ and $0 \leq \theta = \theta(\xi) < 1$ be obtained from (K.6) with ξ satisfying (3.7), where η is defined in (3.6). Then, by Hölder's inequality, Jensen's inequality, Corollaries B.1 and B.2, (3.4), and (3.6),

$$\begin{aligned}
& (D_n - 1)E \left[\frac{\{\mathcal{S}(\hat{k}_{n,\delta_n}) - \mathcal{S}(k_{n,D_n}^*)\}^2}{L_{n,D_n}(k_{n,D_n}^*)} \right] \\
& \leq (D_n - 1) \left[E \frac{\{\mathcal{S}(\hat{k}_{n,\delta_n}) - \mathcal{S}(k_{n,D_n}^*)\}^{2q}}{(L_{n,D_n}(\hat{k}_{n,\delta_n}))^q} \right]^{1/q} \left[E \left\{ \frac{L_{n,D_n}(\hat{k}_{n,\delta_n})}{L_{n,D_n}(k_{n,D_n}^*)} \right\}^{q/(q-1)} \right]^{(q-1)/q} \\
& = O \left(\frac{D_n - 1}{(k_{n,D_n}^*)^{1-\theta(1+\frac{1}{q})}} \right) + o \left(\frac{D_n - 1}{\log \delta_n^{-1}} \right) + O \left(\frac{D_n - 1}{(\log \delta_n^{-1})^{\frac{1}{2}} (k_{n,D_n}^*)^{\frac{1}{2}-\frac{\theta}{q}}} \right) = o(1). \quad (\text{B.49})
\end{aligned}$$

By Corollaries B.1 and B.3 and an argument similar to that used to prove (B.49),

$$(D_n - 1)E \left[\frac{\{\mathbf{f}(\hat{k}_{n,\delta_n}) - \mathbf{f}(k_{n,D_n}^*)\}^2}{L_{n,D_n}(k_{n,D_n}^*)} \right] = o(1). \quad (\text{B.50})$$

Consequently, the desired result is ensured by (B.49), (B.50), and Proposition 2. \square

PROOF OF THEOREM 2. First note that when $\lim_{n \rightarrow \infty} \delta_n = 1$ and condition (i) (or (ii)) of Theorem 2 are assumed instead of (3.4) and (3.5), the left-hand sides of (B.33)-(B.35) still converge to 0. Therefore, (B.29) follows. Let $0 < \xi < (1/2) - \xi_2$ if condition (i) of Theorem 2 holds, and $0 < \xi < \min\{(1/2) - \xi_2, (\delta_1^*/2) - \xi_2\}$ if condition (ii) of Theorem 2 holds. Then, by Jensen's inequality and the same reasoning used in the proofs of Corollaries B.2 and B.3, we have for any $q > 0$,

$$E \left| \frac{\mathcal{S}(\hat{k}_{n,\delta_n}) - \mathcal{S}(k_{n,D_n}^*)}{(L_{n,D_n}(\hat{k}_{n,\delta_n}))^{1/2}} \right|^{2q} = o(1) \text{ and } E \left| \frac{\mathbf{f}(\hat{k}_{n,\delta_n}) - \mathbf{f}(k_{n,D_n}^*)}{(L_{n,D_n}(\hat{k}_{n,\delta_n}))^{1/2}} \right|^{2q} = o(1). \quad (\text{B.51})$$

Consequently, the claimed result follows from (B.29), (B.51), (2.11), and Proposition 2. \square

Appendix C: Proof of Theorem 3

Instead of verifying Theorem 3 directly, we will first investigate the prediction performances of $S_n^{(P_n)}(k)$, defined in (2.15). By an analogy with (4.1) of Shibata (1980),

$$\begin{aligned}
S_n^{(P_n)}(k) &= NL_{n,D_n}(k) + P_n k \left(\hat{\sigma}_n^2(k) - \sigma^2 \right) + \left(k\sigma^2 - N \|\hat{\mathbf{a}}_n(k) - \mathbf{a}(k)\|_{\hat{R}_n(k)}^2 \right) \\
&\quad + N\sigma^2 + N \left(S_{K_n, n-1}^2(k) - \sigma_k^2 \right), \quad (\text{C.1})
\end{aligned}$$

where $D_n = P_n$, for a $k \times k$ symmetric matrix A and a k -dimensional vector \mathbf{y} , $\|\mathbf{y}\|_A = \mathbf{y}'A\mathbf{y}$, and the definition of $S_{K_n, n-1}^2(k)$ can be found in Lemma B.1. Note that the relation, $D_n = P_n$, will be used throughout this appendix. Based on (C.1) and an argument similar to that used in (5.34) of Ing and Wei (2005), we have

$$P(\hat{k}_{n, P_n}^S = k) \leq \sum_{i=1}^5 P(|U_{in}(k)| \geq (1/5) V_{n, D_n}(k)), \quad (\text{C.2})$$

where $V_{n, D_n}(k)$ is defined after (B.2), $\hat{k}_{n, P_n}^S = \arg \min_{1 \leq k \leq K_n} S_n^{(P_n)}(k)$,

$$\begin{aligned} NL_{n, D_n}(k)|U_{1, n}(k)| &= |P_n k(\hat{\sigma}_n^2(k) - \sigma^2)|, \\ NL_{n, D_n}(k)|U_{2, n}(k)| &= |P_n k_{n, D_n}^*(\hat{\sigma}_n^2(k_{n, D_n}^*) - \sigma^2)|, \\ NL_{n, D_n}(k)|U_{3, n}(k)| &= |k\sigma^2 - N\|\hat{\mathbf{a}}_n(k) - \mathbf{a}(k)\|_{\hat{R}_n(k)}^2|, \\ NL_{n, D_n}(k)|U_{4, n}(k)| &= |k_{n, D_n}^*\sigma^2 - N\|\hat{\mathbf{a}}_n(k_{n, D_n}^*) - \mathbf{a}(k_{n, D_n}^*)\|_{\hat{R}_n(k_{n, D_n}^*)}^2|, \text{ and} \\ NL_{n, D_n}(k)|U_{5, n}(k)| &= |S_{K_n, n-1}^2(k) - \sigma_k^2 - S_{K_n, n-1}^2(k_{n, D_n}^*) - \sigma_{k_{n, D_n}^*}^2|. \end{aligned}$$

Theorem C.1. *Let the assumptions of Theorem 3 hold. Then, (2.8) and (2.9) hold for $\hat{k}_{n, OS} = \hat{k}_{n, P_n}^S$, and (4.4) holds with \hat{k}_{n, P_n} replaced by \hat{k}_{n, P_n}^S .*

PROOF. By Lemma B.1 and analogies with Ing and Wei (2005, (5.43) and (5.47)), we have for $q > 0$, all $1 \leq k \leq K_n$, and all sufficiently large n , $E|U_{1, n}(k)|^q \leq C(P_n^q k^q N^{-q} + N^{-q/2})$, $E|U_{2, n}(k)|^q \leq C(P_n^q k_{n, D_n}^{*q} N^{-q} + N^{-q/2})$, $E|U_{3, n}(k)|^q \leq C(k^q N^{-q/2} + k^{q/2} N^{-q} L_{n, D_n}^{-q}(k))$, $E|U_{4, n}(k)|^q \leq C(k_{n, D_n}^{*q} N^{-q/2} + k_{n, D_n}^{*q/2} N^{-q} L_{n, D_n}^q(k))$, and $E|U_{5, n}(k)|^q \leq C\|\mathbf{a}(k) - \mathbf{a}(k_{n, D_n}^*)\|_R^q N^{-q/2} L_{n, D_n}^{-q}(k)$. These moment bounds and an argument similar to that used to verify Corollary B.1 give for $q > 0$,

$$\lim_{n \rightarrow \infty} E \left(\frac{L_{n, D_n}(\hat{k}_{n, P_n}^S)}{L_{n, D_n}(k_{n, D_n}^*)} - 1 \right)^q = 0. \quad (\text{C.3})$$

With the help of (K.1)-(K.6), (4.1), (4.2), and the above moment properties, we can follow the ideas used in the proofs of Corollaries B.2 and B.3 to obtain that for $q > 0$,

$$\begin{aligned} E \left| \frac{\mathcal{S}(\hat{k}_{n, P_n}^S) - \mathcal{S}(k_{n, D_n}^*)}{(L_{n, D_n}(\hat{k}_{n, P_n}^S))^{1/2}} \right|^{2q} &= O((k_{n, D_n}^*)^{-(1-\theta)q+\theta}) + o((P_n - 1)^{-q}) \\ &+ O((P_n - 1)^{-q/2} (k_{n, D_n}^*)^{(-q/2)+\theta}), \end{aligned} \quad (\text{C.4})$$

where $0 \leq \theta = \theta(\xi) < 1$ is obtained from (K.6) with ξ satisfying (4.3), and

$$\lim_{n \rightarrow \infty} E \left| \frac{\mathbf{f}(\hat{k}_{n, P_n}^S) - \mathbf{f}(k_{n, D_n}^*)}{(L_{n, D_n}(\hat{k}_{n, P_n}^S))^{1/2}} \right|^{2q} = o((P_n - 1)^{-q}). \quad (\text{C.5})$$

Consequently, the claimed result is guaranteed by (3.6) (with ξ satisfying (4.3)), (C.3)-(C.5), and Proposition 2. \square

PROOF OF THEOREM 3. It suffices to show that (C.3)-(C.5) hold with \hat{k}_{n,P_n}^S replaced by \hat{k}_{n,P_n} . Define $G_n(k) = N \exp\{\text{IC}_{P_n}(k)\} - S_n^{(P_n)}(k)$ and $|U_{6,n}(k)| = |G_n(k) - G_n(k_{n,D_n}^*)|/NL_{n,D_n}(k)$. Then, by the same reasoning as in (C.2), $P(\hat{k}_{n,P_n} = k) \leq \sum_{i=1}^6 P(|U_{in}(k)| \geq (1/6)V_{n,D_n}(k))$. Moreover, Taylor's expansion and Ing and Wei (2005, (5.42)) yield that for $q > 0$, all $1 \leq k \leq K_n$, and all sufficiently large n , $E|U_{6,n}(k)|^q \leq CP_n^{2q}K_n^{2q}N^{-2q}L_{n,D_n}^{-q}(k)$. These inequalities and the same argument used in the proof of Theorem C.1 give the desired results. \square

Appendix D: Proofs of Theorems 4 and 6

PROOF OF THEOREM 4. First observe that for all sufficiently large n ,

$$p_0 \leq K_n, k_n^* = k_{n,H_n}^* = p_0, L_n(k_n^*) = \frac{p_0\sigma^2}{N}, L_{n,H_n}(k_{n,H_n}^*) = \frac{(H_n - 1)p_0\sigma^2}{N}, \quad (\text{D.1})$$

where $H_n = o(n)$ and $H_n > 1$. Define $I_{3,n}(k) = \{N(\mathbf{f}(k) - \mathbf{f}(p_0))^2\}/(p_0\sigma^2)$. In view of (D.1) and by Hölder's inequality, we have, for all sufficiently large n ,

$$\begin{aligned} (D_n - 1) \frac{E(\mathbf{f}(\hat{k}_{n,\delta_n}) - \mathbf{f}(k_{n,D_n}^*))^2}{L_{n,D_n}(k_{n,D_n}^*)} &\leq \sum_{k=1}^{p_0-1} (E|I_{3,n}(k)|^r)^{\frac{1}{r}} P^{\frac{r-1}{r}}(\hat{k}_{n,\delta_n} = k) \\ + \sum_{k=p_0+1}^{K_n} (E|I_{3,n}(k)|^r)^{\frac{1}{r}} P^{\frac{r-1}{r}}(\hat{k}_{n,\delta_n} = k) &\equiv (I) + (II), \end{aligned} \quad (\text{D.2})$$

where $r > 1$ and $D_n = D_{APE_{\delta_n}}$. According to Ing and Wei (2005, Proposition 1 and Lemmas 1 and 2), (B.7), and (B.11), it can be shown that

$$E|I_{3,n}(k)|^r \leq \begin{cases} CN^r, & 1 \leq k \leq p_0 - 1, \\ Ck^r, & p_0 + 1 \leq k \leq K_n, \end{cases} \quad (\text{D.3})$$

as n is sufficiently large. Armed with (D.3), Lemmas B.2-B.4, the conditions imposed on δ_n , and the fact that for $1 \leq k \leq K_n$ and $k \neq p_0$, $V_{n,D_n}^{-1}(k) \leq C$, the proof of Corollary B.1 is modified to obtain that for any $s > 0$, (I) = $O(n^{-s})$ and (II) = $o((\log \delta_n^{-1})^{-s})$. Hence, (2.9) holds for $\hat{k}_{n,OS} = \hat{k}_{n,\delta_n}$. Similarly, it can be shown that \hat{k}_{n,δ_n} also satisfies (2.8). In view of Proposition 2 and (D.1), (2.3) ((2.4)) is achieved by \hat{k}_{n,δ_n} . Using the above argument and moment bounds for $|U_{in}(k)|$, $i = 1, \dots, 6$ (see Appendix C), (2.8), (2.9), and (2.3) ((2.4)) can also be verified for \hat{k}_{n,P_n} , where P_n satisfies $P_n \rightarrow \infty$ and $P_n/n \rightarrow 0$. The details are omitted in order to save space. \square

PROOF OF THEOREM 6. Unlike the previous theorems, Proposition 2 is not applied in the proof of Theorem 6 since the penalty term associated with $\hat{k}_n^{(\iota)}$, $2I_{\{\hat{k}_{n,P_n} \neq \hat{k}_{n^\iota, P_{n^\iota}}\}} + P_n I_{\{\hat{k}_{n,P_n} = \hat{k}_{n^\iota, P_{n^\iota}}\}}$, is random. In the following, we shall directly verify that

$$\limsup_{n \rightarrow \infty} \frac{q_n(\hat{k}_n^{(\iota)}) - \sigma^2}{L_n(k_n^*)} \leq 1. \quad (\text{D.4})$$

First assume that condition (ii) of Theorem 6 holds. Let $0 < \xi < \min\{\delta_1^*/2, 1/2\}$. Then, there are $0 \leq \theta = \theta(\xi) < 1$ and $M = M(\xi) > 0$ such that (5.2) is fulfilled. Define

$$B_{n,M^*} = A_{P_n, \theta, M}^C \cap \{k : 1 \leq k \leq K_n, \frac{L_{n,P_n}(k) - L_{n,P_n}(k_{n,P_n}^*)}{L_{n,P_n}(k_{n,P_n}^*)} < M^*\},$$

where M^* is some positive constant, and $I_{4,n}(k) = (\mathbf{f}(k) + \mathcal{S}(k))^2 / L_n(k_n^*)$. Then,

$$\begin{aligned} \frac{q_n(\hat{k}_n^{(\iota)}) - \sigma^2}{L_n(k_n^*)} &= E\{I_{4,n}(\hat{k}_n^{(\iota)})\} \leq E\{I_{4,n}(\hat{k}_{n,2})\} \\ &+ E\{I_{4,n}(\hat{k}_{n,P_n})I_{\{\hat{k}_{n,P_n} = \hat{k}_{n^\iota, P_{n^\iota}}\}}(I_{\{\hat{k}_{n,P_n} \in B_{n,M^*}\}} + I_{\{\hat{k}_{n,P_n} \notin B_{n,M^*}\}})\} \equiv (I) + (II). \end{aligned} \quad (\text{D.5})$$

Observe that for $r > 1$,

$$\begin{aligned} (II) &\leq E\{I_{4,n}(\hat{k}_{n^\iota, P_{n^\iota}})I_{\{\hat{k}_{n^\iota, P_{n^\iota}} \in B_{n,M^*}\}}\} + E\{I_{4,n}(\hat{k}_{n,P_n})I_{\{\hat{k}_{n,P_n} \notin B_{n,M^*}\}}\} \\ &= \sum_{\substack{k=1 \\ k \in B_{n,M^*}}}^{K_{n^\iota}} E\{I_{4,n}(k)I_{\{\hat{k}_{n^\iota, P_{n^\iota}} = k\}}\} + \sum_{\substack{k=1 \\ k \notin B_{n,M^*}}}^{K_n} E\{I_{4,n}(k)I_{\{\hat{k}_{n,P_n} = k\}}\} \\ &\leq C \left\{ \sum_{\substack{k=1 \\ k \in B_{n,M^*}}}^{K_{n^\iota}} \frac{L_n(k)}{L_n(k_n^*)} P^{(r-1)/r}(\hat{k}_{n^\iota, P_{n^\iota}} = k) + \sum_{\substack{k=1 \\ k \notin B_{n,M^*}}}^{K_n} \frac{L_n(k)}{L_n(k_n^*)} P^{(r-1)/r}(\hat{k}_{n,P_n} = k) \right\} \\ &\equiv C\{(III) + (IV)\}, \end{aligned} \quad (\text{D.6})$$

where the second inequality follows from Hölder's inequality and the fact that for all $1 \leq k \leq K_n$, $E|\mathbf{f}(k) + \mathcal{S}(k)|^{2r} \leq CL_n^r(k)$, which is ensured by Wei (1987, Lemma 2), Ing and Wei (2005, Proposition 1 and Lemmas 1 and 2), and (B.16). To deal with (III), by (5.3) and the definition of B_{n,M^*} , we have, for all sufficiently large n ,

$$B_{n,M^*} \cap \{1, 2, \dots, K_{n^\iota}\} \subseteq A_{P_{n^\iota}, \theta, M}.$$

Hence, (5.2) ensures that for all $k \in B_{n,M^*} \cap \{1, 2, \dots, K_{n^\iota}\}$ and sufficiently large n ,

$$V_{n^\iota, P_{n^\iota}}^{-1}(k) = \{L_{n^\iota, P_{n^\iota}}(k)\} \{L_{n^\iota, P_{n^\iota}}(k) - L_{n^\iota, P_{n^\iota}}(k_{n^\iota, P_{n^\iota}}^*)\}^{-1} \leq C(k_{n^\iota, P_{n^\iota}}^*)^\xi. \quad (\text{D.7})$$

The definition of B_{n,M^*} also yields for all $k \in B_{n,M^*}$ and $P_n \geq 2$, $L_n(k)/L_n(k_n^*) \leq (P_n - 1)\{L_{n,P_n}(k)/L_{n,P_n}(k_{n,P_n}^*)\} \leq CP_n$. According to this, (D.7), and the moment bounds for $|U_{in}(k)|, i = 1, \dots, 6$, we have, for $q > 0$ and all sufficiently large n ,

$$(III) \leq CP_n(k_{n^\iota, P_{n^\iota}}^*)^{\xi q} \left\{ \sum_{\substack{k=1 \\ k \in B_{n,M^*}}}^{K_{n^\iota}} \frac{P_{n^\iota}^q (k^q + k_{n^\iota, P_{n^\iota}}^{*q})}{N_\iota^q} + \frac{k^q + k_{n^\iota, P_{n^\iota}}^{*q}}{N_\iota^{q/2}} \right. \\ \left. + \frac{k^{q/2} + k_{n^\iota, P_{n^\iota}}^{*q/2}}{N_\iota^q L_{n^\iota, P_{n^\iota}}^q(k)} + \frac{\|\mathbf{a}(k) - \mathbf{a}(k_{n^\iota, P_{n^\iota}}^*)\|_R^q}{N_\iota^{q/2} L_{n^\iota, P_{n^\iota}}^q(k)} + \frac{K_{n^\iota}^{2q} P_{n^\iota}^{2q}}{N_\iota^{2q} L_{n^\iota, P_{n^\iota}}^q(k)} \right\}. \quad (D.8)$$

By taking q on the right-hand side of (D.8) large enough and in view of the restrictions on ξ (given at the beginning of this proof), ι , and P_n ,

$$(III) = o(1). \quad (D.9)$$

Similarly, (5.2) and the definition of B_{n,M^*} imply that for all sufficiently large n and $k \in \{k : 1 \leq k \leq K_n \text{ and } k \notin B_{n,M^*}\}$, (D.7) is still valid if n^ι and P_{n^ι} are replaced by n and P_n , respectively. This finding, the fact that for $P_n \geq 2$, $L_n(k)/L_n(k_n^*) \leq (P_n - 1)\{L_{n,P_n}(k)/L_{n,P_n}(k_{n,P_n}^*)\}$, and an argument similar to the one used to verify (D.9) give (IV) = $o(1)$, which, together with (D.5), (D.6), (D.9), and Theorem 2 of Ing and Wei (2005), yields (D.4).

Next, assume that condition (i) holds. By the moment bounds for $|U_{in}(k)|, i = 1, \dots, 6$, and the similar reasoning to that used in the proof of Theorem 4, we have

$$\lim_{n \rightarrow \infty} P(\hat{k}_{n,P_n} \neq p_0) = 0, \quad (D.10)$$

and for any $q > 0$,

$$E|I_{4,n}(\hat{k}_{n,2})|^q = O(1). \quad (D.11)$$

Since

$$\frac{q_n(\hat{k}_n^{(\iota)}) - \sigma^2}{L_n(k_n^*)} \leq E\{I_{4,n}(\hat{k}_{n,P_n})\} + E\{I_{4,n}(\hat{k}_{n,2})(I_{\{\hat{k}_{n,P_n} \neq p_0\}} + I_{\{\hat{k}_{n^\iota, P_{n^\iota}} \neq p_0\}})\}, \quad (D.12)$$

(D.4) follows from (D.10)-(D.12), Hölder's inequality, and $\limsup_{n \rightarrow \infty} E(I_{4,n}(\hat{k}_{n,P_n})) \leq 1$ (which is ensured by Theorem 4). This completes the proof of the theorem. \square

Acknowledgements. I would like to thank a co-editor, an associate editor, and two anonymous referees for their valuable and constructive comments, which helped to

substantially improve the readability and content of the paper. I am also grateful to Atsushi Inoue and Lutz Kilian for kindly correcting some errors in my GAUSS code for implementing the simulations in a previous version of this paper.

REFERENCES

- Akaike, H. (1969). Fitting autoregressive models for prediction. *Ann. Inst. Statist. Math.* **21** 243-247.
- Akaike, H. (1974). A new look at the statistical model identification. *IEEE Trans. Automat. Control* **19** 716-723.
- Bhansali, R. J. (1986). Asymptotically efficient selection of the order by the criterion autoregressive transfer function. *Ann. Statist.* **14** 315-325.
- Findley, D. F. (2005). Asymptotic second moment properties of out-of-sample forecast errors of misspecified regARIMA models and the optimality of GLS. *Statist. Sinica*, **15** 447-476.
- Findley, D. F. and Wei, C. Z. (1993). Moment bounds for deriving time series CLT's and model selection procedures. *Statist. Sinica* **3** 453-470.
- Goldenshluger, A. and Zeevi, A. (2001). Nonasymptotic bounds for autoregressive time series modeling. *Ann. Statist.* **29** 417-444.
- Hannan, E. J., McDougall, A. J. and Poskitt, D. S. (1989). Recursive estimation of autoregressions. *J. R. Statist. Soc. B* **51** 217-233.
- Hannan, E. J. and Quinn, B. C. (1979). The determination of the order of an autoregression. *J. R. Statist. Soc. B* **41** 190-195.
- Hasen, M. H. and Yu, B. (2001). Model selection and the principle of minimum description length. *J. Am. Statist. Assoc.* **96** 746-774.
- Hemerly, E. M. and Davis, M. H. A. (1989). Strong consistency of the PLS criterion for order determination of autoregressive processes. *Ann. Statist.* **17** 941-946.
- Hurvich, C. M. and Tsai, C. L. (1989). Regression and time series model selection in small samples. *Biometrika* **76** 297-307.
- Ing, C. K. (2001). A note on mean-squared prediction errors of the least squares predictors in random walk models. *J. Time Ser. Anal.* **22** 711-724.
- Ing, C. K. (2003). Multistep prediction in autoregressive processes. *Economet. Theory* **19** 254-279.
- Ing, C. K. (2005). Accumulated prediction errors, information criteria, and optimal forecasting for autoregressive time series. Technical Report. Available at <http://www.stat.sinica.edu.tw/Ing/newaic3.pdf>.
- Ing, C. K. and Wei, C. Z. (2003). On same-realization prediction in an infinite-order autoregressive process. *J. Multivariate Anal.* **85** 130-155.

- Ing, C. K. and Wei, C. Z. (2005). Order selection for same-realization prediction in autoregressive processes. *Ann. Statist.* **33** 2423-2474.
- Inoue, A. and Kilian, L. (2006). On the selection of forecasting models. *J. Economet.* **130** 273-306.
- Karagrigoriou, A. (1997). Asymptotic efficiency of the order selection of a nongaussian AR process. *Statist. Sinica* **7** 407-423.
- Kavalieris, L. (1989). The estimation of the order of an autoregression using recursive residuals and cross-validation. *J. Time Ser. Anal.* **10** 271-281.
- Kunitomo, N. and Yamamoto, T. (1985). Properties of predictors in misspecified autoregressive time series. *J. Am. Statist. Assoc.* **80** 941-950.
- Lai, T. L. and Wei, C. Z. (1982). Least squares estimates in stochastic regression models with application to identification and control systems. *Ann. Statist.* **10** 154-166.
- McCracken, M. W. (2000). Robust out-of-sample inference. *J. Economet.* **99** 195-223.
- Mallows, C. L. (1973). Some comments on C_p . *Technometrics* **12** 591-612.
- Rissanen, J. (1986). Order estimation by accumulated prediction errors. In *Essays in Time Series and Allied Processes* (J. Gani and M. P. Priestley, eds.) *J. Appl. Probab.* **23A** 55-61.
- Schorfheide, F. (2005). VAR forecasting under misspecification. *J. Economet.* **128** 99-136.
- Schwarz, G. (1978). Estimating the dimension of a model. *Ann. Statist.* **6** 461-464.
- Shibata, R. (1976). Selection of the order of an autoregressive model by Akaike's information criterion. *Biometrika* **63** 117-126.
- Shibata, R. (1980). Asymptotically efficient selection of the order of the model for estimating parameters of a linear process. *Ann. Statist.* **8** 147-164.
- Speed, T. P. and Yu, B. (1993). Model selection and prediction : normal regression. *Ann. Inst. Statist. Math.* **45** 35-54.
- Wax, M. (1988). Order selection for AR models by predictive least squares," *IEEE Trans. Acoust. Speech Signal Process.* **36** 581-588.
- Wei, C. Z. (1987). Adaptive prediction by least squares predictors in stochastic regression models with application to time series. *Ann. Statist.* **15**, 1667-1682.
- Wei, C. Z. (1992). On predictive least squares principles. *Ann. Statist.* **20**, 1-42.
- West, K. D. (1996). Asymptotic inference about predictive ability. *Econometrica* **64** 1067-1084.
- Zheng, X. and Loh, W. Y. (1997). A consistent variable selection criterion for linear models with high-dimensional covariates. *Statist. Sinica* **7** 311-325.