

Econometrics 710  
Final Exam, Spring 2011

The exam consists of one question, broken in several parts.  
The model is

$$y_i = \mathbf{x}_i' \boldsymbol{\beta} + e_i \quad (1)$$

$$E(\mathbf{z}_i e_i) = 0 \quad (2)$$

The dimensions are:  $\mathbf{x}_i$ ,  $\mathbf{z}_i$ , and  $\boldsymbol{\beta}$  are  $k \times 1$ ,  $k > 1$ , and  $y_i$  and  $e_i$  are  $1 \times 1$ . Let

$$\mathbf{Q} = \begin{bmatrix} \mathbf{Q}_{xx} & \mathbf{Q}_{xz} \\ \mathbf{Q}_{zx} & \mathbf{Q}_{zz} \end{bmatrix} = \begin{bmatrix} E(\mathbf{x}_i \mathbf{x}_i') & E(\mathbf{x}_i \mathbf{z}_i') \\ E(\mathbf{z}_i \mathbf{x}_i') & E(\mathbf{z}_i \mathbf{z}_i') \end{bmatrix}$$

Assume both  $\mathbf{Q}_{xx}$  and  $\mathbf{Q}_{xz}$  have full rank  $k$ .

Let  $\hat{\boldsymbol{\beta}}$  be the least-squares estimate obtained by regressing  $y_i$  on  $\mathbf{x}_i$ , and let  $\tilde{\boldsymbol{\beta}}$  be the 2SLS estimator obtained by estimation of (1) using the instrument  $\mathbf{z}_i$ .

1. Find

$$\delta = \text{plim}_{n \rightarrow \infty} (\hat{\boldsymbol{\beta}} - \tilde{\boldsymbol{\beta}})$$

2. Suppose that in addition to (1) and (2),

$$E(\mathbf{x}_i e_i) = 0 \quad (3)$$

Quite simply, what does this condition mean? What is  $\delta$  under this assumption?

3. Write the difference  $\hat{\boldsymbol{\beta}} - \tilde{\boldsymbol{\beta}}$  as a function of sample moments of  $\mathbf{x}_i$ ,  $\mathbf{z}_i$ , and  $e_i$ .
4. Under (1)-(3), find the asymptotic distribution of

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n \begin{pmatrix} \mathbf{x}_i \\ \mathbf{z}_i \end{pmatrix} e_i$$

as  $n \rightarrow \infty$ .

5. Under (1)-(3) find the asymptotic distribution of  $\sqrt{n} (\hat{\boldsymbol{\beta}} - \tilde{\boldsymbol{\beta}})$  as  $n \rightarrow \infty$ .

6. Suppose that

$$E(e_i^2 | \mathbf{x}_i, \mathbf{z}_i) = \sigma^2 \quad (4)$$

How does the asymptotic variance from question 5 simplify under (4)?

7. Propose an estimator of the asymptotic variance under (4).
8. Propose a test statistic for (3) under (4) and find its asymptotic distribution under the assumption that  $\mathbf{Q} > 0$
9. Describe how to use this statistic to test the hypothesis that  $\mathbf{x}_i$  is exogenous.
10. Extra Credit. Show where  $\mathbf{Q} > 0$  is used in the answer to question 8.